

Unnecessary Deaths: The Human and Financial Costs of Hospital Infections

By
Betsy McCaughey, Ph.D
Chairman
Committee to Reduce Infection Deaths
www.hospitalinfection.org



Essential Facts You Need to Know About Hospital Infections

Keep in mind:

- We have the knowledge to prevent hospital infection deaths.
- We don't have to wait for a scientific breakthrough.
- Yet most hospitals have failed to act.
- The situation is growing more dangerous because, increasingly, hospital infections cannot be cured with commonly used antibiotics

1. Infections contracted in hospitals are the fourth largest killer in America. Every year in this country, two million patients¹ contract infections in hospitals, and an estimated 103,000 die as a result,² as many deaths as from AIDS, breast cancer, and auto accidents combined.³
2. A few hospitals in the U.S. are proving that infections are almost entirely preventable. How are they doing it? Through rigorous hand hygiene, meticulous cleaning of equipment and rooms in between patient use, testing incoming patients to identify those carrying dangerous bacteria and strictly isolating those who test positive, and requiring caregivers to use gowns or disposable aprons when treating positive patients.
3. In 2003, the Society for Healthcare Epidemiologists of America (SHEA) announced the precautions that, research proves, can eradicate most hospital infections. Yet only a few hospitals have taken these precautions, and the CDC still has not called on all hospitals to implement them.
4. Hospital infections add an estimated \$30 billion to the nation's hospital costs each year. Patients, insurers and taxpayers pay part of that cost, but hospitals have to absorb much of the cost. As a result, infections erode hospital profits. Preventing infections can turn a financially failing hospital profitable.
5. Better infection control in hospitals is essential to prepare the nation for avian flu or bioterrorism. If avian flu were to wing its way to the U.S., the death toll would depend largely on what American hospitals did when the first avian flu patients were admitted. If hospitals have effective infection controls in place, they can prevent bird flu from infecting other patients who did not come in with it. If not, bird flu could sweep through hospitals. Right now, most hospitals are woefully underprepared. Hospitals have failed to stop the spread of ordinary infections spread by touch and would not be able to contain flu viruses, which are communicated by droplets from coughing and sneezing as well as touch. Even more challenging would be smallpox, plague, and other bioterrorism weapons that can travel through the air. Shoddy infection control is poor preparation for a flu epidemic and poor homeland security as well.
6. Hospital infection is a far deadlier problem than the number of uninsured. The Institute of Medicine estimates that as many as 18,000 people a year die prematurely because they don't have health insurance. That's tragic. But five times as many people die each year from hospital infections, and most of them are insured.⁴

Dear Reader,

This report has been written to help you and to enlist your help in correcting a deadly situation that kills an estimated 103,000 people in this country each year—as many deaths as from AIDS, breast cancer, and auto accidents combined.⁵

Where does it kill? In our hospitals. What is it? Hospital infection.

The death toll is staggering. So is the economic cost. Hospital infections add over \$30 billion a year to what the nation spends on hospital care, enough to pay for a large part of the Medicare prescription drug program.⁶

These infections are almost all preventable. A few hospitals in the U.S. are proving it, reducing some of the deadliest types of infections by 90 percent. Their achievements prove that we have the knowledge to solve this problem. No major scientific breakthrough is needed. What is lacking is leadership.

That is why I founded the Committee to Reduce Infection Deaths (RID): to motivate hospitals to make infection prevention a top priority; to inform patients about the steps they can take to reduce their risk of infection; and to ensure that no matter where you live, you can find out which hospitals in your area have the worst infection problems.

■ **RID holds forums** for hospital administrators, public health officials, lawmakers, medical educators, insurers, and patient advocates, showing them how infections can be eradicated and *how much money can be saved*. The humanistic reasons to stop hospital infections are obvious. RID forums also make a compelling economic case for infection control, showing that for some hospitals, preventing infection can actually make the difference between profitability and loss.

■ **RID educates** the public through television, radio, popular publications, and our website. One of our most important educational tools is the “14 Steps You Can Take to Reduce the Risk of a Hospital Infection,” which is included in this study.

■ **RID works with state lawmakers and other policy makers** to develop hospital infection report cards, because if you need to be hospitalized, you should be able to choose a hospital with a low infection rate.

■ **RID partners with health insurers** to develop incentives for hospitals to improve infection control and to deliver life-saving information to patients.

■ **RID is encouraging medical schools and nursing schools** to educate their students about how bacteria are spread from patient to patient in hospitals and the precautions that should be taken to protect their patients—a subject that is almost entirely neglected in most schools.

The Committee has a distinguished membership, including Dr. Sherwin Nuland, author of *The Doctor's Plague*, a biography of Ignac Semmelweis; Dr. Elizabeth Whelan, founder of the American Council on Science and Health, and Nobel Laureate Dr. Joshua Lederberg, as well as corporate leaders, philanthropists, and civic leaders. Our goal is to save lives.

Everyday you hear about health care problems such as providing for the uninsured. The Institute of Medicine recently estimated that as many as 18,000 people a year may die prematurely because they don't have health insurance.⁷ But consider this even more tragic fact: Five times that many people die each year from hospital infection, and most of them are insured. Having insurance is no guarantee that you will be safe in the hospital. The only way to ensure that is to clean up this deadly problem.

Betsy McCaughey, Ph.D.
Founder and Chairman,
Committee to Reduce Infection Deaths
www.hospitalinfection.org



About the National Center for Policy Analysis

The National Center for Policy Analysis (NCPA) is a nonprofit, nonpartisan public policy research institute established in 1983. Its mission is to seek innovative private sector solutions to public policy problems. The NCPA focuses on issues of importance nationally and internationally. It is headquartered in Dallas, Texas, in order to draw on “outside-the-beltway” thinking, and maintains an active office in Washington, D.C., in order to inform deliberations in Congress and the Executive Branch.

The NCPA is probably best known for developing the concept of Medical Savings Accounts (MSAs), now known as Health Savings Accounts (HSAs). In 2003, as part of Medicare reform, Congress and the President made HSAs available to all nonseniors, potentially revolutionizing the entire health care industry. The NCPA home page (www.ncpa.org) links visitors to the best available information, including studies produced by think tanks all over the world.

Table of Contents

Third World Hygiene in Our First Class Medical System. 1

The Major Problem: Poor Hygiene 3

Infections Erode Hospital Profits 5

The Importance of Hospital Infection Report Cards 7

Shouldn't Medical Students Be Taught Hygiene? 10

Success Stories: Infections *Can* Be Eradicated 11

 A. Dr. Carlene Muto Describes Victory Over MRSA at
 The University Of Pittsburgh Medical Center 11

 B. Dr. Richard Shannon Aims for Zero Infections at
 Allegheny Hospital 12

 C. Dr. Barry Farr Recalls Early Victories at the
 University Of Virginia Hospital. 14

14 Steps You Can Take to Reduce Your Risk of a Hospital Infection 16

Appendix A:
RID's Model Hospital Infection Report Card Bill. 19

Appendix B:
Society for Healthcare Epidemiologists of America (SHEA)
Guideline for Preventing Nosocomial Transmission of
Multidrug-Resistant Strains of *Staphylococcus aureus*
and *Enterococcus* 21

Appendix C:
The Institute for Healthcare Improvement's
100,000 Lives Campaign 44

About The Author. 46

Endnotes 47

I

Third World Hygiene in Our First Class Medical System

Every day in hospitals across the United States wondrous medical procedures rescue patients from the brink of death. But there's a catch. In these same hospitals, hygiene is so inadequate that one out of every twenty patients contracts an infection.⁸

Infections that have been nearly eradicated in some countries are raging through American hospitals. In 2003, the Society for Healthcare Epidemiologists of America warned that although hospitals have infection control programs, "there is little evidence of control in most facilities."⁹

The danger is worsening because hospital infections, increasingly, cannot be tamed with commonly used antibiotics. One of the deadliest germs is *methicillin-resistant Staphylococcus aureus* (or MRSA). Patients who do survive MRSA often spend months in the hospital and endure repeated surgeries to cut out infected tissue. In 1974, 2 percent of *Staph* infections were MRSA. By 1995, the number had climbed to 22%, in 2003 an alarming 57% and still rising.¹⁰

Denmark, Holland, and Finland once faced similar rates, but brought them down below 1 percent.¹¹ How? Through rigorous hand hygiene, meticulous cleaning of equipment and rooms in between patient use, and testing incoming hospital patients to identify those carrying MRSA and strictly isolating them to prevent transmission to other patients. Wheelchairs and other equipment used to transport patients who test positive for MRSA are not used for other patients, and hospital staff have to change their uniforms and footwear after entering the rooms of MRSA patients, before they are permitted in other areas of the hospital.

A few hospitals—very few—in the United States are proving these precautions work here too. The University of Virginia Hospital eradicated MRSA.¹² The Veterans Hospital in Pittsburgh reduced MRSA by 85 percent in a pilot program.¹³ The University of Pittsburgh-Medical Center-Presbyterian Hospital slashed MRSA by 90 percent in the medical intensive care units.¹⁴ Twenty-nine healthcare

institutions in Iowa eliminated another drug-resistant germ, *vancomycin-resistant Enterococcus* (or VRE).¹⁵ Unfortunately, most U.S. hospitals have not implemented these precautions. Here's what you'll find in the great unwashed majority of hospitals.

II

The Major Problem: Poor Hygiene

Over half the time, physicians and other caregivers break the most fundamental rule of hygiene by failing to clean their hands before treating patients.¹⁶ Gloves are not the solution, because pulling them on with dirty hands contaminates the gloves.

Nearly three quarters of patients' rooms are contaminated with MRSA and VRE. These bacteria are on cabinets, counter tops, over-the-bed tables, bed rails, and other surfaces.¹⁷ Once patients and caregivers touch these surfaces, their hands become *vectors for disease*. Ordinary cleaning solutions are effective against these bugs, but surfaces need to be drenched for several minutes, not just sprayed and wiped quickly.¹⁸

Astoundingly, most hospitals in the United States don't routinely test patients to determine which ones are carrying MRSA and other bacteria. Seventy to ninety percent of patients carrying MRSA are never identified. Knowing which patients are sources of infection is key to stopping the spread.¹⁹

Clothing is frequently a conveyor belt for infections. When doctors and nurses lean over a patient with

MRSA, the white coats and uniforms pick up bacteria 65 percent of the time and carry it to other patients.²⁰ Hospitals that are conquering infection require their staff to put on fresh gowns or disposable aprons every time they treat patients with MRSA. (The aprons cost a nickel and are ripped off rolls like clear plastic dry-cleaning bags.)

A major academic hospital in New York City is struggling to control the spread of *Clostridium difficile*, an infection usually caused by fecal material from one patient entering another patient's mouth. How could that happen? Doctors there suspect it's because clinical nursing assistants wear the same clothes while doing two jobs: emptying bed pans and delivering food trays.²¹

The privacy curtains that surround a patient's bed are seldom changed, though they are often the last thing a caregiver touches before treating a patient, and the first thing touched afterward, when the caregiver reaches up with contaminated gloves to pull open the curtain. Stethoscopes, blood pressure monitors, and other pieces of equipment are frequently carrying live bacteria.

Does your doctor clean the stethoscope before listening to your chest? Probably not, though the American Medical Association recommends it.²²

We have the knowledge to prevent infection. What has been lacking is the will. Most hospitals have not made fighting infection a top priority. The federal Centers for Disease Control and Prevention are also to blame. The CDC has tracked the rapid increase in drug-resistant hospital infections for a quarter century but has not advocated the rigorous steps necessary to stop them.

In 2003, the Society for Healthcare Epidemiologists of America urged hospitals to routinely test incoming patients, isolate those who test positive for MRSA and VRE, and prevent

the spread of these bacteria to other patients via equipment, staff clothing, and room surfaces. "There are at least fifty studies demonstrating the effectiveness of these precautions," explains Dr. Carlene Muto, an epidemiologist at the University of Pittsburgh Medical Center, and "not one study suggesting it's possible to control MRSA without them."²³ One study shows that MRSA spreads from patient to patient 15 times as fast under CDC standards as under the more rigorous precautions advocated by the Society.²⁴

Yet two years later, the CDC has still not called on hospitals to implement these precautions. Hospitals must be happy at the lack of pressure, but the public should be outraged. Every year of delay is costing thousands of lives.

III


Infections Erode Hospital Profits

■

Many hospital administrators say they can't afford to take these precautions. The truth is, they can't afford *not* to. Infections erode hospital profits, because rarely are hospitals fully paid for the added weeks or months patients must spend in the hospital when they get an infection. For example: Pennsylvania's Allegheny Hospital would have made a profit treating a 37-year old video programmer and father of four who was admitted with acute pancreatitis, but the economics changed when the patient developed a MRSA bloodstream infection. He had to stay in the hospital 86 days, and the hospital lost \$41,813.²⁵ Pittsburgh's Shadyside Hospital tamed a MRSA outbreak and saved ten dollars for every dollar spent on improving hygiene, testing patients, and isolating those with MRSA.²⁶

Infections add more than \$30 billion annually to the nation's health costs.²⁷ The tab will increase rapidly, as more infections become drug-resistant.²⁸

Additional research on the cost of infections shows that:



■ Post surgical wound infections more than double a patient's hospital costs. When a patient develops an infection after surgery, the cost of care increases 119 percent, on average, at a teaching hospital, and 101 percent at a community hospital.³⁰

■ Urinary tract infections increase a patient's hospital costs by 47 percent at a teaching hospital and 35 percent at a community hospital.³¹

■ The average ventilator-associated pneumonia infection (a type

of infection contracted when a patient is on a respirator) adds **\$40,000** to a patient's hospital costs.³²

■ A central catheter-related bloodstream infection increases a patient's hospital costs by about **\$30,000** on average.³³

■ *Staphylococcus aureus* infections are especially costly. According to a recent nationwide study, patients with *Staph* infections incur hospital costs that amount to more than triple the average hospital costs of other patients.³⁴

If the cost is not enough to motivate hospital leaders to improve infection controls, they now face

two other pressures. One is from the trial lawyers. Remember asbestos? Hospital infection is the next asbestos. The infection problem has all the hot button essentials of a successful class action lawsuit: two million helpless victims a year, copious evidence that infections are preventable, and a consistent pattern of failure to act.

Secondly, hospitals in many states will soon be compelled to come clean about their infection rates. Six states—Florida, Missouri, Pennsylvania, New York, Illinois, and Virginia—recently enacted laws to provide the public with risk-adjusted hospital infection report cards, and several other states are poised to follow.

IV

The Importance of Hospital Infection Report Cards

Maureen Daly wishes she had known more when she took her 63-year old mother to the hospital. Johanna had slipped and broken her shoulder at a restaurant, and no one expected that she would be in the hospital for more than a day or two. But a *Staph* infection ravaged her body for four months and killed her. “What happened to my mother shouldn’t happen to anyone,” says Daly. “If only I had had enough information to choose a hospital with a better infection record.”

If you need to be hospitalized, wouldn’t you want to know which hospital in your area has the lowest infection rate? Good luck getting that information!

Most states don’t even collect data on hospital infections. Twenty-one states require hospitals to report infections serious enough to cause severe injury or death,³⁵ but the requirement is seldom enforced, and worse still, states go along with the hospital industry’s demands to keep the data secret. The federal Centers for Disease Control and Prevention collect infection data from several hundred hospitals around the nation, but the CDC also

promises hospitals to keep infection rates secret.³⁶ Government, for the most part, is not helping you choose a safe hospital.

The irony is that it’s easy to get information for the less important decisions you make in life, such as where to have lunch. Most states will help you find out which restaurants and delicatessens have been cited for health violations. But you can’t find out which hospital has the worst infection rate. You can go home to make your own sandwich, but you can’t perform surgery on yourself.

The good news is that Florida, Missouri, Pennsylvania, Illinois, Virginia, and New York recently passed laws to provide the public with hospital infection report cards. Publicly comparing hospital performance will motivate hospitals to improve.

New York’s experience with another type of hospital report card proves this. In 1989, New York became the first state to publish each hospital’s risk-adjusted mortality rate for cardiac bypass surgery. The results? Deaths from bypass surgery dropped 40 percent, giving

New York the lowest mortality rate in the nation for that procedure.³⁷ Critics of hospital report cards speculate that deaths went down in New York because hospitals avoided treating the sickest patients, fearing that high-risk operations would bring down the hospital's grade. However, the evidence proves that's untrue. Deaths declined for a different reason: hospitals forced their worst-performing surgeons—generally, those with low volume—to stop doing the procedure. Patients of the 27 barred surgeons were more than three times as likely to die during surgery. In technical jargon, the 27 surgeons had an average risk-adjusted mortality rate of 11.9 percent, compared with a statewide average of 3.1 percent.³⁸ Wisconsin also found that report cards motivate poorly performing hospitals to improve, according to a 2001 study of 24 hospitals there.³⁹

Is there a reason *not* to have infection report cards? The hospital industry argues that publicly comparing hospital infection rates would be unfair to hospitals that treat AIDS, cancer, and organ transplant patients who are especially vulnerable to infection. Fair enough, but reports can be risk-adjusted to reflect these differences. What is unfair is keeping the public uninformed.

Fortunately, 30 states are considering legislation to provide the public with the information they need. These states should use the model bill suggested here (Appendix A), because it improves upon the laws already passed in

three ways: First, it specifies the method of risk-adjustment for surgical site infections used by the CDC, rather than leaving the risk-adjustment method to be determined by committee. This should assure hospitals that comparisons will be fair and take into account which hospitals treat especially sick and infection-prone patients.⁴⁰

Secondly, the bill imposes civil penalties on hospitals that fail to report or flagrantly underreport their infections. These penalties are needed. For many years, some hospitals have openly ignored data collection laws with impunity. For example, in one recent year, hospitals in New York reported only 16.5 percent of the post-surgical deaths that the law required them to report.⁴¹ In 2005, the first year of Pennsylvania's hospital infection reporting program, hospitals reported only one tenth as many infections to the new program as they billed. Some Pennsylvania hospitals implausibly claimed they had no infections at all.⁴²

Thirdly, the model bill ensures that hospital infection reporting will benefit the public, not enrich trial lawyers. The bill provides that "none of the data collected and reported under this law can be used in litigation against an individual hospital."

Next time you hear an ad on the radio urging you to use a particular hospital because it has the best doctors or the latest equipment, keep in mind what you're *not* being told: how many patients get infections

while in that hospital. Hospitals are doing their best to keep that information secret. In contrast, in England hospital infection rates are posted conspicuously on the front door of the hospital. Americans deserve the same information. The

legislation proposed here won't help hospitals save face, but it will help you choose a safe hospital. Let hospitals vie for your business by improving their infection rates.

V

Shouldn't Medical Students Be Taught Hygiene?

What else needs to be done? Medical schools should be teaching future doctors the precautions they must take to protect their patients from infection. It's hard to believe, but most medical schools devote virtually no time, not even one full class, to showing students how bacteria are transmitted from patient to patient on clothing, equipment, and gloves, and what specifically they should be doing to prevent it. Dr. Frank Lowey, a professor at the New York-Presbyterian Hospital at the Columbia University Medical Center says, "it's something we should have done quite a while ago." Lowey says it's ironic that "there are curriculum committees devoted to making sure that bioterrorism is covered, and the risk of nosocomial infections far outweighs that."⁴³

Some medical schools are stressing the importance of curbing the use of antibiotics.⁴⁴ That's good, because overuse of antibiotics wastes money and causes bacteria to morph into new, drug-resistant strains. But limiting the use of antibiotics won't stop hospital infections. Patients who contract MRSA get it from unclean hands or contaminated equipment or clothing, not simply from taking antibiotics. No hospital has ever eradicated infection merely by controlling the use of these drugs.

When medical students put on their white coats and swear the Hippocratic Oath, they should be taught *how* to do no harm. Preventing the spread of bacteria is an essential part of that lesson. They should learn it before they go out on the hospital floors and touch their first patient.

VI

Success Stories: Infections *Can Be Eradicated*



A. Dr. Carlene Muto Describes Victory Over MSRA at the University of Pittsburgh Medical Center⁴⁵

“It’s a fabulous feeling,” says Dr. Carlene Muto, reflecting on the team effort that has resulted in a 90 percent reduction in *methicillin-resistant Staphylococcus aureus* (MRSA) in the medical intensive care unit at the hospital where she is director of infection control. How long did it take? Three years. Ask her how it was done. She explains that it required total commitment from the top leadership at the hospital and caregivers.

When Muto came to UPMC-Presbyterian, the flagship hospital in the University of Pittsburgh system, in the 1990s, drug-resistant *Staphylococcus aureus* was a rapidly growing problem. In 2000, Muto launched a campaign to eradicate the “superbug” in the hospital’s medical intensive care unit. Critical to the strategy was active surveillance culturing—meaning that every patient coming into the intensive care unit who might be carrying MRSA was cultured. Muto, one of the co-authors of *The Society for Healthcare Epidemiologists of*

America’s guideline, emphasizes that you can’t eliminate infection until you know which patients are the sources of the bacteria. Every patient who tested positive was isolated, and doctors and nurses treating them wore gowns and masks, and kept equipment used on these patients away from others. By 2003, MRSA was almost eliminated. The strategy has worked so well that it has now been expanded to all 15 intensive care units in the hospital system.

The key, explains Muto, was to identify every patient carrying the dangerous bacteria. “We had total compliance, 98 percent to 100 percent, with culturing patients,” she said, adding that she was astonished. After all, asking nurses to culture every new patient in the ICU meant adding one more thing to an already long list of tasks they have to do. The staff reaction, says Muto, “has been overwhelmingly positive.” “That’s essential,” she adds. “You can come up with an idea, but no matter how great it is, you have

to have the buy in from the staff at the point of care.”

Getting caregivers to clean their hands has been a tougher challenge, in part because at the beginning, Muto explains, some “nurses didn’t realize that if they went into a room of a patient in isolation and didn’t touch the patient or the bed linen but did touch other surfaces such as countertops, their hands *were* contaminated.”

Now that the education process is well under way, hand cleaning compliance is about 69 percent, well above the national average but not good enough for Muto and her team. The top leadership

at UPMC-Presbyterian is taking an uncompromising position on the failure of staff and doctors to clean their hands. The hospital is getting set to impose stiff penalties, including firing staff members who chronically ignore hand cleaning rules and denying doctors the privilege of practicing at the hospital.

The goal? “Our goal is 100 percent compliance with hand cleaning, 100 percent compliance with gowning, 100 percent compliance with surveillance culturing,” says Muto, adding excitedly that she can only imagine what can be achieved when they reach perfection.



B. Dr. Richard Shannon Aims for Zero Infections

When Dr. Richard Shannon told the top executives at Allegheny Hospital that he wanted to do something about central line-associated blood stream infections (CLABs), the hospital leadership expected him to suggest reducing them by 10 or 20 percent over several years. To their surprise, Shannon said he wanted to totally eradicate these deadly infections in ninety days. And he did it! Even more amazing, he and his staff kept these infections near zero in the medical intensive care unit and coronary care unit during the entire next year, achieving a 95 percent reduction in CLAB-related deaths.⁴⁶

Why strive for merely minor improvement when lives are at stake? Shannon’s pet peeve is

benchmarking—the thinking all too common in hospitals today that it’s okay to have infections and medical errors so long as they don’t exceed the national average. “Who volunteers to have a family member get one of the infections we plan on having this year?” The goal has to be zero infections and perfect care, says Shannon, who is Chairman of the Department of Medicine at Allegheny.⁴⁷

How was that goal reached? By ensuring that all caregivers meticulously follow a regimen for inserting and removing central lines that includes masks, gowns, gloves, and drapes; inserting lines in the neck area rather than in the groin area, which is more difficult to keep

clean; rearranging supply closets to ensure that the supplies needed to carry out this regimen are easily accessible, even when staff are rushed; and empowering all staff members to enforce hand cleaning and other rules of hygiene. If a doctor doesn't clean his hands, the nurse working alongside can call a halt to the procedure until the doctor complies.

Shannon oversees some 800 employees and a \$150 million budget. Nevertheless, he makes time to speak across the country, with PowerPoint in tow, showing his audiences that preventing infections is possible *and profitable*. Doing the right thing costs less, he says, using Allegheny's financial records to prove the point. A typical example is the tragic case of a woman who came into the hospital for stomach reduction surgery, a procedure that should have produced a \$9,900 gross profit for the hospital. But when she developed a central line-associated bloodstream infection and had to spend 47 days in the hospital, that profit turned into a \$16,000 loss. Preventing CLABs saved Allegheny \$1.4 million the first year.⁴⁸

The best news of all is that the success at Allegheny is being duplicated by at least a few other institutions. At Johns Hopkins, catheter-related bloodstream infections in the intensive care unit have been virtually eliminated. How?

ICU staff were educated about the seriousness of catheter-related infections; a catheter-insertion cart was created to ensure that necessary equipment was readily at hand; doctors were asked daily whether catheters should be removed; bedside nurses were given a safety checklist to follow during insertion; and nurses were empowered to stop procedures if safety rules were not being followed. Peter Provonost, the intensive care physician at Johns Hopkins who developed the safety checklist, sees the success as proof that infections are not inevitable.⁴⁹

That is Richard Shannon's mantra as well. Shannon is amazed that so little is being done nationwide to curb bloodstream infections *and* to halt the alarming rise in MRSA. Shannon asks why the procedures that reduced *Staph* infections by 85 percent in a pilot program at the V.A. Hospital in Pittsburgh are not being duplicated everywhere. "What if you had a patient with TB or SARS? Wouldn't you pull out all the stops, gloving and gowning and washing up all the time? Well, we haven't seen TB in years, and we've never seen SARS, but we have MRSA silently stalking us every day." The magnitude of the problem, he says, is "a call to action for all health-care providers to step up and get serious about all hospital-acquired infections."⁵⁰



C. Dr. Barry Farr Recalls Early Victories at the University of Virginia Hospital⁵¹

Barry Farr remembers the first outbreak of MRSA at the University of Virginia Hospital. It was 1978, and Farr and his wife had recently come to the hospital to train, having just finished medical school. “MRSA was wildly out of control,” he recalls, and the hospital was doing “what most American healthcare facilities are still doing today.” As a result, the hospital “failed miserably to control the MRSA.”

For nearly three years, as the outbreak raged on, the hospital followed a business-as-usual approach: no routine cultures were being taken to identify the patients silently carrying the bacteria. The result, recalls Farr, was that doctors were touching patients who had MRSA, or allowing their white coats to brush up against them, and then passing the bacteria on to other patients without knowing it. At the hospital infection control meetings, the mood was pessimistic and apathetic. Staff members were saying “no one has ever controlled this.”

Finally, after three years of failure, the hospital took a radical step, inspired by the success of several European countries that had brought MRSA under control. The hospital began regularly testing patients for the bacteria and isolating those who tested positive. The results were stunning. Soon after the testing began, in December 1980, MRSA declined rapidly, and

by the summer of 1982, the hospital was MRSA free. “It was beautiful,” Farr recalls.

Surveillance culturing—identifying every patient carrying the bacteria—was the key to thwarting the outbreak and eradicating MRSA, says Farr. It was to work again a decade later.

The University of Virginia hospital was struck with MRSA a second time in the early 1990s, when a surgeon apparently walked into the neonatal intensive care unit with MRSA on his hands or clothing and transmitted it to one of the babies. Quickly it was spread to babies in the neighboring bassinets, and then to another neonatal intensive care unit when one of the babies carrying the bacteria was moved there. The hospital immediately put into place the same precautions that had worked a decade earlier, and the outbreak was curtailed. Culturing every baby, and isolating everyone who tested positive, was once again the key.

Would this method conquer other deadly bacteria as well? Soon afterward, the hospital faced an outbreak of *vancomycin-resistant Enterococci* (VRE), which spread rapidly to 30 percent of patients on eight separate wards. After several months, the hospital brought the outbreak under control once again by testing patients, isolating the carriers, and making sure that all staff

put on gowns and gloves when treating them.

Are the University of Virginia's successes atypical? "No," says Farr. "There are over ninety studies, probably 100 by now," demonstrating that this method works. Yet antibiotic-resistant infections are "clearly out of control in the American health care setting." Why? Farr suggests that faulty cost-cutting is partly to blame.

Hospital administrators complain about the cost of these rigorous precautions, but the data proves these precautions save money. Farr compared the University of Virginia hospital with several other university hospitals of similar size. These other hospitals "are spending between \$1 million and \$3 million a year extra to treat antibiotic-resistant infec-

tions, far more than what UVA has had to spend on gowns, cultures, and gloves. We're taking the ounce of prevention approach. Many other hospitals are taking the pound of cure approach."

Another reason few hospitals are adopting rigorous infection control is that the public has not demanded it. "In Britain there is a public outcry over the failure to control MRSA infections in hospitals, and the British government is reportedly now considering firing hospital directors that fail to take effective measures to control MRSA," says Farr. "In this country there has been comparatively little outcry from the public and no urgent demands from the government that the spread of infections be stopped."

14 Steps You Can Take to Reduce Your Risk of a Hospital Infection



1. Ask that hospital staff clean their hands before treating you. This is the single most important way to protect yourself in the hospital. If you're worried about being too aggressive, just remember your life could be at stake. All caregivers should clean their hands before treating you. Alcohol-based hand cleaners are more effective at removing most bacteria than soap and water.⁵² Do not hesitate to say the following to your doctor or caregiver: "Excuse me, but there's an alcohol dispenser right there. Would you mind using that before you touch me, so I can see it?" Don't be falsely assured by gloves. Gloves more often protect staff than patients. If caregivers have pulled on gloves without cleaning their hands first, the gloves are already contaminated before they touch you.⁵³

2. Before your doctor uses a stethoscope to listen to your chest, ask that the diaphragm (or flat surface of the stethoscope) be wiped with alcohol. Numerous studies show that stethoscopes are often contaminated with *Staphylococcus*

aureus and other dangerous bacteria, because caregivers seldom take the time to clean them in between patient use.⁵⁴ The American Medical Association recommends that stethoscopes routinely be cleaned for each patient. The same precautions should be taken for many other commonly used pieces of equipment too.

3. Ask visitors to clean their hands and avoid sitting on your bed.⁵⁵

4. If you need surgery, choose a surgeon with a low infection rate. Surgeons know their rate of infection for various procedures. Ask for it. If they won't tell you, consider choosing another surgeon. You should be able to compare hospital infection rates too, but that information is almost impossible to get. That is why RID is working hard for hospital infection report cards in every state.

5. Beginning one week before surgery, shower frequently with Chlorhexidine soap. Various brands

can be found at drug stores. This will help remove any dangerous bacteria you may be carrying on your own skin.

6. Ask your surgeon to have you tested for *Staphylococcus aureus* at least one week before you come into the hospital. The test is simple, usually just a nasal swab. About one third of people carry *Staphylococcus aureus* on their skin, and if you are one of them, extra precautions can be taken to protect you from infection, to give you the correct antibiotic during surgery, and to prevent you from transmitting bacteria to others.

7. On the day of your operation, remind your doctor that you may need an antibiotic one hour before the first incision. For many types of surgery, a pre-surgical antibiotic is the standard of care, but it is often overlooked by busy hospital staff.⁵⁶

8. Ask your doctor about keeping you warm during surgery. Operating rooms are often kept cold for the comfort of the staff, but research shows that for many types of surgery, patients who are kept warm resist infection better.⁵⁷ There are many ways to keep patients warm, including special blankets, hats and booties, and warmed IV liquids.

9. Do not shave the surgical site. Razors can create small nicks in the skin, through which bacteria

can enter. If hair must be removed before surgery, ask that clippers be used instead of a razor.⁵⁸

10. Ask that your surgeon limit the number of personnel (including medical students) in the operating room. Every increase in the number of people adds to your risk of infection.⁵⁹

11. Ask your doctor about monitoring your glucose (sugar) levels continuously during and after surgery, especially if you are having cardiac surgery. The stress of surgery often makes glucose levels spike erratically. New research shows that when blood glucose levels are tightly controlled to stay between 80–110 mg/unit, heart patients resist infection better. Continue monitoring even when you are discharged from the hospital, because you are not fully healed yet.⁶⁰

12. Avoid a urinary tract catheter if possible. It is a common cause of infection. The tube allows urine to flow from your bladder out of your body. Sometimes catheters are used when busy hospital staff don't have time to walk patients to the bathroom. Ask for a diaper or bed pan instead. They're safer.⁶¹

13. If you must have an IV, make sure that it is inserted and removed under clean conditions and changed every 3 to 4 days. Intravenous catheters, or IVs, are a common source of infection and are not always necessary. If you

need one, insist that it be inserted and removed under clean conditions, which means that your skin is cleaned at the site of insertion, and the person treating you is wearing clean gloves. Alert hospital staff immediately if any redness appears.

14. If you are planning to have your baby by Cesarean section, take the steps listed above as if you were having any other type of surgery.

Appendix A

RID's Model Hospital Infection Report Card Bill



The following outline is intended to help state lawmakers as they draft legislation to provide the public with hospital infection rates:

AN ACT to provide the public with information on infection rates at hospitals in the state of _____.

Section 1. Definitions.

(a) The public health law is amended to add a new section (lawmakers here should include the specific title of the public health or health department law to be amended).

(b) "Hospital" shall mean (lawmakers here should consider whether to include only acute care hospitals or also free-standing outpatient surgical centers).

(c) "Hospital-acquired infection" shall mean, as defined by the federal Centers for Disease Control and Prevention (CDC), "any localized or systemic condition resulting from an adverse reaction to the presence of an infectious agent(s) or its toxin(s) that (a) occurs in a patient in a hospital, (b) and was found not to be present or incubating at the time of admission to the hospital, unless (c) the infection was related to a previous admission to the same hospital."

(d) "Risk adjustment" shall mean a statistical procedure for comparing patient outcomes, taking into account the differences in patient populations, including risk factors such as the number of patients on central line catheters, or the number of patients undergoing specific types of surgery, as a percentage of the overall number of patients treated. For purposes of this bill, risk adjustment shall duplicate the CDC's NNIS System surgical wound infection risk index or use the number of central-catheter days as a risk-adjustment factor for central line infections.

Section 2.

(a) Using established public health surveillance methods, each hospital shall maintain a program of identifying and tracking the following types of hospital-acquired infections for the purpose of reporting such data semi-annually to the state health department (lawmakers insert the appropriate state department

Appendix A

here): central line-associated, laboratory confirmed primary bloodstream infections contracted by intensive care unit patients, and surgical site infections.⁶²

(b) The state health department (lawmakers insert the appropriate department name here) shall establish an advisory committee that includes recognized experts in the field of hospital-acquired infection, public reporting of hospital data, and health care quality management to establish data collection and analysis methodologies and risk adjustment procedures.

(c) The state health department (lawmakers insert the appropriate department name here) shall establish a state-wide database of all risk-adjusted, hospital-specific infection rates and make it available to the public on a website and in printed materials that can be used by consumers, purchasers of healthcare, and advocacy groups to compare the performance of individual hospitals, and the aggregate performance of hospitals in the state with those in other states and nationwide.

(d) The first year of data submission under this section shall be considered the “pilot phase” of the reporting system. The pilot phase is to ensure the completeness and accuracy of hospital reporting and the fairness and completeness of the state health department’s report to the public. During this pilot phase, hospital identifiers shall be encrypted, the state health department (lawmakers insert proper department name here) shall provide each hospital with an encryption key for that hospital only, and no public hospital comparisons will be available. Sixty days after the end of the second year of data submission, the state health department (appropriate department name here) will provide its first report to the public with hospital-specific infection rates included.

(e) To ensure compliance with this law and the accuracy of self-reporting by the hospitals, the department shall establish an audit process. A civil penalty of \$_____ shall be imposed on any hospital that fails to report on time, or is shown to substantially underreport infections, for each semi-annual reporting period.

(f) None of the data collected and reported under this law can be used in litigation against an individual hospital.

Appendix B

Society for Healthcare Epidemiologists of America (SHEA) Guideline for Preventing Nosocomial Transmission of Multidrug-Resistant Strains of *Staphylococcus aureus* and *Enterococcus*



Strength of Recommendations

Category Type	Category Subtype	Recommendation
I	A	Strongly recommended for implementation and strongly supported by well-designed experimental, clinical, or epidemiologic studies.
	B	Strongly recommended for implementation and supported by some experimental, clinical, or epidemiologic studies and a strong theoretical rationale.
	C	Required for implementation, as mandated by federal regulation, state regulation, or both or standard.
II		Suggested for implementation and supported by suggestive clinical or epidemiologic studies or a theoretical rationale.
No recommendation		Unresolved issue. Practices for which insufficient evidence or no consensus regarding efficacy exists.

Recommendations

I. Active Surveillance Cultures to Identify the Reservoir for Spread

1. Implement a program of active surveillance cultures and contact precautions to control the spread of epidemiologically significant antibiotic-resistant pathogens known to be spreading in the healthcare system via direct and indirect contact. (IA)29,30,43,45-47,49,57,96,99,102,106,119,138-147,149,171-173,176
2. Surveillance cultures are indicated at the time of hospital admission for patients at high risk for carriage of MRSA, VRE, or both. (IB)71,76,177,320,321
3. Periodic (eg, weekly) surveillance cultures are indicated for patients remaining in the hospital at high risk for carriage of MRSA, VRE, or both because of ward location, antibiotic therapy, underlying disease, duration of stay, or all four. (IA)30,57,102,137,141,147-149,174,181

Appendix B

4. In facilities found to have a high prevalence on initial sampling, a facility-wide culture survey is indicated to identify all colonized patients and allow implementation of contact precautions. (IB)102,145,322
5. Because transmission occurs throughout the healthcare system, these measures should be implemented in all types of healthcare facilities throughout the system. (IB)119,161,176,182,323
6. The frequency of active surveillance cultures should be based on the prevalence of the pathogen and risk factors for colonization. For example, more frequent cultures are needed in a facility where 50% of all *S. aureus* isolates are MRSA than in one where less than 1% of all *S. aureus* isolates are MRSA. (IB)29,30,43,45-47,49,57,96,99,102,106,119,138-147,149,171-173,176
7. The goal of this program should be to identify every colonized patient, so that all colonized patients are cared for in contact (or cohort) isolation to minimize spread to other patients. (IB)29,30,43,45-47,49,57,96,99,102,106,119,138-147,149,171-173,176
8. Surveillance cultures for VRE should use stool samples or swab samples from the rectum or perirectal area. Polymerase chain reaction, culture with broth enhancement, and quantitative stool culture have each been more sensitive than directly plated rectal or perirectal swab cultures, but the latter have been associated with control of infections and can be recommended as effective and cost-effective until less costly methods of using the other procedures become available. (IB)99,102,106,137,149,181
9. VRE patients can be routinely cohorted with other VRE patients. (ID)102,106,145
10. Surveillance cultures for MRSA should always include samples from the anterior vestibule of the nose. (IB)78,315,324
11. If present, areas of skin breakdown should also be sampled for MRSA. (IB)315,324
12. Throat cultures have been shown to detect *S. aureus* and MRSA with sensitivity equal to or greater than that of nasal cultures in multiple patient populations. If used, the throat swab can be plated onto the same agar as the nasal swab. This would enhance sensitivity without adding the cost of an extra culture. (IB)67,74
13. Perirectal–perineal cultures have been shown to detect MRSA with high sensitivity in certain patient populations, but the perirectal–perineal area should not be selected as the only site for culture. (IB)315,324,325

Appendix B

14. Patients colonized or infected with MRSA isolates can be cohorted with other MRSA patients. (II)30,43,45

15. Patients with MRSA isolates that are eradicable because of known susceptibility to multiple drugs useful for eradication (eg, mupirocin, rifampin, minocycline, trimethoprim-sulfamethoxazole, or all four) should not be cohorted with those with isolates resistant to these drugs, if eradication will be used as an adjunctive measure. (II)272

16. In certain settings, such as nursing homes and psychiatric wards, identification of colonized patients is important, but contact precautions may require modification allowing for social contact while limiting physical contact. (II)119,182,323

II. Hand Hygiene

1. HCWs should be encouraged to decontaminate (clean) their hands with an antiseptic-containing preparation before and after all patient contacts. (IA)121,326-330

2. Soap and water hand washing is required when hands are visibly dirty or visibly contaminated with blood, body fluids, or body substances. (IA)331

3. When hands are not visibly contaminated with blood, body fluids, or body substances, use of an alcohol hand rub containing an emollient should be encouraged. (IB)215,332-338

4. Lotion compatible with (ie, that does not inactivate) the antiseptic being used should be provided for use by HCWs. (II)339-343

5. Monitoring of hand hygiene compliance and feedback to HCWs should be done to motivate greater compliance. (IB)215,344

III. Barrier Precautions for Patients Known or Suspected to Be Colonized or Infected With Epidemiologically Important Antibiotic-Resistant Pathogens Such as MRSA or VRE

1. Gloves should always be worn to enter the room of a patient on contact precautions for colonization or infection with antibiotic-resistant pathogens such as MRSA, VRE, VISA, or VRSA. (IA)122,132,212,225-230

2. Gowns always should be worn as part of contact precautions for all patient and environmental contact with patients known to be colonized by antibiotic-resistant pathogens such as MRSA, VRE, VISA, or VRSA, except when there is no direct contact with patient or environmental surfaces. (IA)29,30,43,45,47,49,57,59,96,99,102,106,119,122,132,135,136,138-147,149,171-173,176,345

Appendix B

3. Universal gown and glove use or universal gloving alone also can be considered for adjunctive control on high-risk wards among patients with surveillance cultures pending. (IB)37,44,105,316-318,346

4. Masks should be worn as part of isolation precautions when entering the room of a patient colonized or infected with MRSA, VISA, or VRSA to decrease nasal acquisition by HCWs. (II)30,123,124,129,231,232

IV. Antibiotic Stewardship

1. Avoid inappropriate or excessive antibiotic prophylaxis and therapy. (IB)194,251,347

2. Ensure correct dosage and duration of antibiotic therapy. (IB)348-350

3. Restrict the use of vancomycin (if possible and appropriate for care of the individual patient being treated) to decrease the selective pressure favoring vancomycin resistance. (IB)115,269

4. To prevent the establishment of VRE intestinal colonization, decrease the use of agents with little or no activity against enterococci, such as third-generation and fourth-generation cephalosporins, in patients not known to be VRE colonized (if possible and appropriate for care of the individual patient being treated). (IB)115,267,268,351,352

5. To prevent persistent high-density VRE colonization, decrease the use of antianaerobic agents in patients with known VRE intestinal colonization (if possible and appropriate for care of the individual patient being treated). (II)102,113,159,270

6. To help prevent persistent carriage of MRSA, reduce the use of antibiotics and particularly fluoroquinolones to the minimum necessary in institutions where MRSA is endemic. (IB)251-258

7. Avoid therapy for colonization except when suppression or eradication of colonization is being attempted using an evidence-based approach for infection prevention, for psychological benefit of the patient, or for cost benefit (ie, by reducing the need for long-term isolation). (IB)5,272,285,286

V. Decolonization or Suppression of Colonized Patients

1. Consider MRSA decolonization therapy for both patients and HCWs as an adjunctive measure for controlling spread of MRSA in selected populations when appropriate. (IB)30,176,271,272,275-277

2. Any program of decolonization therapy should incorporate routine susceptibility testing, as selection of inactive agents is less likely to achieve eradication. (II)272,353

Appendix B

3. Widespread use, prolonged use, or both of decolonization therapy should be avoided, because this has been associated with the evolution and spread of antibiotic-resistant strains, undermining the effectiveness of the control effort. (IB)285,286

VI. Other

1. Educational programs should be conducted to ensure that HCWs understand why antibiotic-resistant pathogens are epidemiologically important, why prevention of spread is critically necessary for control, and which measures for preventing spread have proven effective. (IB)215,220

2. Ensure that the hospital method of disinfecting hospital surfaces for antibiotic-resistant organisms (especially VRE) has been shown to be adequate based on the results of studies of such methods in the healthcare setting or perform cultures in the room of discharged patients to confirm the adequacy of terminal cleaning. This requires review of the disinfectant agent, method and meticulousness of cleaning, dilutions, and contact time. (IB)102,161,169,294

3. Use the hospital computer system to record longterm isolation indicators for patients colonized with MRSA, VRE, VISA, or VRSA so that on return the computer will provide an alert regarding the need for isolation. (IB)297

4. Dedicate the use of noncritical patient-care equipment to a single patient (or cohort of patients infected or colonized with the pathogen requiring precautions) to avoid sharing between patients. If use of common equipment or items is unavoidable, then adequately clean and disinfect them before use for another patient. (IB)99,150-155,296

References

- Centers for Disease Control and Prevention. *Campaign to Prevent Antimicrobial Resistance in Healthcare Settings: Why a Campaign?* Atlanta, GA: Centers for Disease Control and Prevention; 2001. Available at www.cdc.gov/drugresistance/healthcare/problem.htm.
- Neu HC. The crisis in antibiotic resistance. *Science* 1992;257:1064-1073.
- Marshall G, Crofton JW, Cruickshank R, et al. The treatment of pulmonary tuberculosis with isoniazid. *BMJ* 1952;2:735-746.
- Aubry-Damon H, Soussy CJ, Courvalin P. Characterization of mutations in the rpoB gene that confer rifampin resistance in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1998;42:2590-2594.
- Schmitz FJ, Fluit AC, Hafner D, et al. Development of resistance to ciprofloxacin, rifampin, and mupirocin in methicillin-susceptible and -resistant *Staphylococcus aureus* isolates. *Antimicrob Agents* Vol. 24 No. 5 PREVENTING SPREAD OF ANTIBIOTIC RESISTANCE 379 *Chemother* 2000;44:3229-3231.
- O'Neill AJ, Cove JH, Chopra I. Mutation frequencies for resistance to fusidic acid and rifampin in *Staphylococcus aureus*. *J Antimicrob Chemother* 2001;47:647-650.
- Eisenstadt E, Carlton BC, Brown BJ. Gene mutation. In: Gerhardt P, Murray RGE, Wood WA, Krieg NR, eds. *Methods for General and Molecular Bacteriology*. Washington, DC:

Appendix B

- American Society for Microbiology; 1994:297-316.
- 8 Kucers A, Bennett N. *The Use of Antibiotics*, 4th ed. London: Heinemann Medical Books; 1987.
 - 9 Holmes O. The contagiousness of puerperal fever. *N Engl Q J Med Surg* 1842-1843;1:501-530.
 - 10 Semmelweis I. *The Etiology, the Concept and the Prophylaxis of Childbed Fever*. Pest, Hungary: CA Hartleben's Verlag-Expedition; 1861.
 - 11 Darwin C. *On the Origin of Species by Means of Natural Selection*. London: J. Murry; 1859.
 - 12 Centers for Disease Control and Prevention. National Nosocomial Infections Surveillance (NNIS) System report: data summary from January 1990–May 1999. *Am J Infect Control* 1999;27:520-532.
 - 13 Centers for Disease Control and Prevention. *NNIS Antimicrobial Resistance Report: Vancomycin-Resistant Enterococcus (VRE) Facts*. Atlanta, GA: Centers for Disease Control and Prevention; 1999.
 - 14 Centers for Disease Control and Prevention. National Nosocomial Infection Surveillance (NNIS) System report: data summary from January 1992–June 2001. *Am J Infect Control* 2001;29:404-421.
 - 15 Cosgrove SE, Sakoulas G, Perencevich EN, Schwaber MJ, Karchmer AW, Carmeli Y. Comparison of mortality associated with methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* bacteremia: a meta-analysis. *Clin Infect Dis* 2003;36:53-59.
 - 16 Carmeli Y, Cosgrove SE, Harbarth S, Karchmer AW, Kaye KS, Qi Y. The impact of methicillin-resistance in *Staphylococcus aureus* bacteremia on patient outcomes: mortality, length of stay and hospital charge. Presented at the 41st Annual Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; December 16-19, 2001; Chicago, IL. Abstract K-1221:415.
 - 17 Engemann JJ, Carmeli Y, Cosgrove SE, et al. Adverse clinical and economic outcomes attributable to methicillin resistance among patients with *Staphylococcus aureus* surgical site infection. *Clin Infect Dis* 2003;36:592-598.
 - 18 Salgado C, Farr B. The cost of vancomycin-resistance (VR): a metaanalysis. Presented at the 12th Annual Meeting of the Society for Healthcare Epidemiology of America; April 6-9, 2002; Salt Lake City, UT. Abstract 113:67.
 - 19 Benson K, Hartz AJ. A comparison of observational studies and randomized controlled trials. *N Engl J Med* 2000;342:1878-1886.
 - 20 Concato J, Shah N, Horwitz RJ. Randomized controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000;342:1887-1892.
 - 21 Hiramatsu K. Molecular evolution of MRSA. *Microbiol Immunol* 1995;39:531-543.
 - 22 Hiramatsu K, Cui L, Kuroda M, Ito T. The emergence and evolution of methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol* 2001;9:486-493.
 - 23 Kreiswirth B, Kornblum J, Arbeit WE, et al. Evidence for a clonal origin of methicillin-resistance in *Staphylococcus aureus*. *Science* 1993;259:227-230.
 - 24 Oliveira DC, Tomasz A, de Lencastre H. The evolution of pandemic clones of methicillin-resistant *Staphylococcus aureus*: identification of two ancestral genetic backgrounds and the associated mec elements. *Microb Drug Resist* 2001;7:349-361.
 - 25 Musser J, Kapur V. Clonal analysis of methicillin-resistant *Staphylococcus aureus* from intercontinental sources: association of the mec gene with divergent phylogenetic lineages implies dissemination by horizontal transfer and recombination. *J Clin Microbiol* 1992;30:2058-2063.
 - 26 Givney R, Vickery A, Holliday A, Pegler M, Benn R. Evolution of an endemic methicillin-resistant *Staphylococcus aureus* population in an Australian hospital from 1967-1996. *J Clin Microbiol* 1998;36:552-556.
 - 27 Crisostomo MI, Westh H, Tomasz A, Chung M, Oliveria DC, de Lencastre H. The evolution of methicillin resistance in *Staphylococcus aureus*: similarity of genetic backgrounds in historically early methicillin-susceptible and resistant and contemporary epidemic clones. *Proc Natl Acad Sci U S A* 2001;98:9865-9870.
 - 28 Enright MC, Robinson DA, Randle G, Feil DJ, Grundmann H, Spratt BG. The evolution-

Appendix B

- ary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc Natl Acad Sci U S A* 2002;99:7687-7692.
- ²⁹ Haley RW, Cushion NB, Tenover FC, et al. Eradication of endemic methicillin-resistant *Staphylococcus aureus* infections from a neonatal intensive care unit. *J Infect Dis* 1995;171:614-624.
- ³⁰ Jernigan JA, Titus MG, Groschel DHM, Getchell-White SI, Farr BM. Effectiveness of contact isolation during a hospital outbreak of methicillin-resistant *Staphylococcus aureus*. *Am J Epidemiol* 1996;143:496-504.
- ³¹ Salmenlinna S, Lyytikäinen O, Kotilainen P, Scotford R, Siren E, Vuopio-Varkila J. Molecular epidemiology of methicillin-resistant *Staphylococcus aureus* in Finland. *Eur J Clin Microbiol Infect Dis* 2000;19:101-107.
- ³² Roberts RB, de Lancastre A, Eisner W, et al. Molecular epidemiology of methicillin-resistant *Staphylococcus aureus* in 12 New York hospitals: MRSA collaborative group. *J Infect Dis* 1998;178:164-171.
- ³³ de Lancastre H, Severina EP, Roberts RB, Kreiswirth B, Tomasz A. Testing the efficacy of a molecular surveillance network: methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus faecium* (VREF) genotypes in six hospitals in the metropolitan New York City area. The BARG Initiative Pilot Study Group. Bacterial Antibiotic Resistance Group. *Microb Drug Resist* 1996;2:343-351.
- ³⁴ Roman RS, Smith J, Walker M, et al. Rapid geographic spread of a methicillin-resistant *Staphylococcus aureus* strain. *Clin Infect Dis* 1997;25:698-705.
- ³⁵ Villari P, Farullo C, Torre I, Nani E. Molecular characterization of methicillin-resistant *Staphylococcus aureus* (MRSA) in a university hospital in Italy. *Eur J Epidemiol* 1998;14:802-816.
- ³⁶ Diekema DJ, Pfaller MA, Turnidge J, et al. Genetic relatedness of multidrug-resistant, methicillin-resistant *Staphylococcus aureus* bloodstream isolates: SENTRY antimicrobial resistance surveillance centers worldwide. *Microb Drug Resist* 2000;6:213-221.
- ³⁷ Vriens MR, Fluit AC, Troelstra A, Verhoef J, Van Der Werken C. Are MRSA more contagious than MSSA in a surgical intensive care unit. *Infect Control Hosp Epidemiol* 2002;23:491-494.
- ³⁸ Witte W, Cuny C, Bräulke C, Heuck D, Klare I. Widespread dissemination of epidemic MRSA in German hospitals. *Eurosurveillance* 1997;2:25-28.
- ³⁹ Austin DJ, Anderson RM. Transmission dynamics of epidemic methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant enterococci in England and Wales. *J Infect Dis* 1999;179:883-891.
- ⁴⁰ Deplano A, Witte W, Van Leeuwen WJ, Brun Y, Struelens MJ. Clonal dissemination of epidemic methicillin-resistant *Staphylococcus aureus* in Belgium and neighboring countries. *Clin Microbiol Infect* 2000;6:239-245.
- ⁴¹ Galdbart JO, Morvan A, El Solh N. Phenotypic and molecular typing of nosocomial methicillin-resistant *Staphylococcus aureus* strains susceptible to gentamicin isolated from France from 1995-1997. *J Clin Microbiol* 2000; 8:185-190.
- ⁴² Sanchez IS, Ramirez M, Troni H, et al. Evidence for geographic spread of a methicillin-resistant *Staphylococcus aureus* clone between Portugal and Spain. *J Clin Microbiol* 1995;33:1234-1236.
- ⁴³ Thompson RL, Cabezudo I, Wenzel RP. Epidemiology of nosocomial infections caused by methicillin-resistant *Staphylococcus aureus*. *Ann Intern Med* 1982;97:309-317.
- ⁴⁴ Verhoef J, Beaujean D, Blok H, et al. A Dutch approach to methicillin-resistant *Staphylococcus aureus*. *Eur J Clin Microbiol Infect Dis* 1999; 18:461-466.
- ⁴⁵ Jernigan JA, Clemence MA, Stott GA, et al. Control of methicillin resistant *Staphylococcus aureus* at a university hospital: one decade later. *Infect Control Hosp Epidemiol* 1995;16:686-696.
- ⁴⁶ Chaix C, Durand-Zaleski I, Alberti C, Brun-Buisson C. Control of endemic methicillin-resistant *Staphylococcus aureus*: a cost benefit analysis in an intensive care unit. *JAMA* 1999;282:1745-1751.
- ⁴⁷ Jans B, Suetens C, Struelens M. Decreasing MRSA rates in Belgian hospitals: results from the national surveillance network after introduction of national guidelines. *Infect*

Appendix B

Control Hosp Epidemiol 2000;21:419.

- 48 Bager F. *DANMAP 98: Consumption of Antimicrobial Agents and Occurrence of Antimicrobials in Bacteria From Food Animals, Food and Humans in Denmark*. Copenhagen: Statens Serum Institut, Danish Veterinary and Food Administration, Danish Medicines Agency, Danish Veterinary Laboratory; 1998. Available at www.svs.dk/dk/z/Danmap%201998.pd. 1999.
- 49 Harbarth S, Martin Y, Rohner P, Henry N, Auckenthaler R, Pittet D. Effect of delayed infection control measures on a hospital outbreak of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 2000;46:43-49.
- 50 Merrer J, Santoli F, Appere de Vecchi C, Tran B, De Jonghe B, Outin H. "Colonization pressure" and risk of acquisition of methicillin-resistant *Staphylococcus aureus* in a medical intensive care unit. *Infect Control Hosp Epidemiol* 2000;21:718-723.
- 51 Farr BM, Salgado CD, Karchmer TB, Sherertz RJ. Can antibiotic-resistant nosocomial infections be controlled? *Lancet Infect Dis* 2001;1:38-45.
- 52 Vincent JL, Bihari DJ, Suter PM, et al. The prevalence of nosocomial 380 INFECTION CONTROL AND HOSPITAL EPIDEMIOLOGY May 2003 infection in intensive care units in Europe: results of the European Prevalence of Infection in Intensive Care (EPIC) Study. EPIC International Advisory Committee. *JAMA* 1995;274:639-644.
- 53 Cars O, Molstad S, Melander A. Variation in antibiotic use in the European Union. *Lancet* 2001;357:1851-1853.
- 54 Frank MO, Batteiger BE, Sorensen SJ, et al. Decrease in expenditures and selected nosocomial infections following implementation of an antimicrobial-prescribing improvement program. *Clinical Performance and Quality Healthcare* 1997;5:180-188.
- 55 Fukatsu K, Saito HK, Matsuda T, Ikeda S, Furukawa S, Muto T. Influences of type and duration of antimicrobial prophylaxis on an outbreak of methicillin-resistant *Staphylococcus aureus* and on the incidence of wound infection. *Arch Surg* 1997;132:1320-1325.
- 56 Batteiger BE. Personal communication. Indianapolis: Indiana University; 2001.
- 57 Back NA, Linnemann CC Jr, Staneck JL, Kotagal UR. Control of methicillin-resistant *Staphylococcus aureus* in a neonatal intensive-care unit: use of intensive microbiologic surveillance and mupirocin. *Infect Control Hosp Epidemiol* 1996;17:227-231.
- 58 Barrett FF, McGehee RF, Finland M. Methicillin-resistant *Staphylococcus aureus* at Boston City Hospital. *N Engl J Med* 1968;279:441-448.
- 59 Boyce JM. Are the epidemiology and microbiology of methicillin-resistant *Staphylococcus aureus* changing? *JAMA* 1998;279:623-624.
- 60 Brumfitt W, Hamilton-Miller J. Methicillin-resistant *Staphylococcus aureus*. *N Engl J Med* 1989;320:1188-1196.
- 61 Naimi TS, LeDell KH, Boxrud D, et al. Epidemiology and clonality of community-acquired methicillin-resistant *Staphylococcus aureus* in Minnesota, 1996-1998. *Clin Infect Dis* 2001;33:990-996.
- 62 Frank AL, Marcinak JF, Mangat PD, Schreckeberger PC. Community-acquired and clindamycin-susceptible methicillin-resistant *Staphylococcus aureus* in children. *Pediatr Infect Dis J* 1999;18:993-1000.
- 63 Suggs AH, Maranan MC, Boyle-Vavra S, Daum RS. Methicillin-resistant and borderline methicillin-resistant asymptomatic *Staphylococcus* colonization in children without identifiable risk factors. *Pediatr Infect Dis J* 1999;18:410-414.
- 64 Herold BD, Immergluck LC, Maranan MC, et al. Community-acquired methicillin in children with no identified predisposing risk. *JAMA* 1998;279:593-598.
- 65 Adecock PM, Pastor P, Medley F, Patterson JE, Murphy TV. Methicillin-resistant *Staphylococcus aureus* (MRSA) in two child care centers. *J Infect Dis* 1998;178:577-580.
- 66 Embil J, Ramotar K, Romance L, et al. Methicillin-resistant *Staphylococcus aureus* in tertiary care institutions on the Canadian prairies 1990-1992. *Infect Control Hosp Epidemiol* 1994;15:646-651.

Appendix B

- ⁶⁷ Shahin R, Johnson IL, Jamieson F, McGeer A, Tolkin J, Ford-Jones EL. Methicillin-resistant *Staphylococcus aureus* carriage in a child care center following a case of disease. *Arch Pediatr Adolesc Med* 1999;153:864-868.
- ⁶⁸ Centers for Disease Control and Prevention. Four pediatric deaths from community-acquired methicillin-resistant *Staphylococcus aureus*: Minnesota and North Dakota, 1997-1999. *MMWR* 1999;48:707-710.
- ⁶⁹ Morin CA, Hadler JL. Population-based incidence and characteristics of community-onset *Staphylococcus aureus* infections with bacteremia in 4 metropolitan Connecticut areas, 1998. *J Infect Dis* 2001;184:1029-1034.
- ⁷⁰ Layton MC, Hierholzer WJ, Patterson JE. The evolving epidemiology of methicillin-resistant *Staphylococcus aureus* at a university hospital. *Infect Control Hosp Epidemiol* 1995;16:12-17.
- ⁷¹ Troillet N, Carmeli Y, Samore MH, et al. Carriage of methicillin-resistant *Staphylococcus aureus* at hospital admission. *Infect Control Hosp Epidemiol* 1998;19:181-185.
- ⁷² Groom AV, Wolsey DH, Naimi TS, et al. Community-acquired methicillin-resistant *Staphylococcus aureus* in a rural American Indian community. *JAMA* 2001;286:1201-1205.
- ⁷³ Salmenlinna S, Lyytikäinen O, Vuopio-Varkila J. Community-acquired methicillin-resistant *Staphylococcus aureus*, Finland. *Emerg Infect Dis* 2002;8:602-607.
- ⁷⁴ Sa-Leao R, Sanches IS, Couto I, Alves CR, de Lencastre H. Low prevalence of methicillin-resistant strains among *Staphylococcus aureus* colonizing young and healthy members of the community in Portugal. *Microb Drug Resist* 2001;7:237-245.
- ⁷⁵ Shopsin B, Mathema B, Martinez J, et al. Prevalence of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* in the community. *J Infect Dis* 2000;182:359-362.
- ⁷⁶ Muto CA, Cage EG, Durbin LJ, Simonton BM, Farr BM. The utility of culturing patients on admission transferred from other health care facilities for methicillin-resistant *Staphylococcus aureus* (MRSA). Presented at the Ninth Annual Meeting of the Society for Healthcare Epidemiology of America; April 18-20, 1999; San Francisco, CA. Abstract M33:67.
- ⁷⁷ Calfee DP, Durbin LJ, Germanson TP, Toney DM, Smith EB, Farr BM. Prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) among household contacts of individuals with nosocomially-acquired MRSA. *Infect Control Hosp Epidemiol*. 2003. In press.
- ⁷⁸ Sanford MD, Widmer AF, Bale MJ, Jones RN, Wenzel RP. Efficient detection and long-term persistence of the carriage of methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis* 1994;19:1123-1128.
- ⁷⁹ Gold HS. Vancomycin-resistant enterococci: mechanisms and clinical observations. *Clin Infect Dis* 2001;33:210-219.
- ⁸⁰ Bonten MJ, Willems R, Weinstein RA. Vancomycin-resistant enterococci: why are they here, and where do they come from? *Lancet Infect Dis* 2001;1:314-325.
- ⁸¹ Murray BE. What can we do about vancomycin-resistant enterococci? *Clin Infect Dis* 1995;20:1134.
- ⁸² Martone WJ. Spread of vancomycin-resistant enterococci: why did it happen in the United States? *Infect Control Hosp Epidemiol* 1998;19:539-545.
- ⁸³ Carmeli Y, Samore MH, Huskins C. The association between antecedent vancomycin treatment and hospital-acquired vancomycin-resistant enterococci: a meta-analysis. *Arch Intern Med* 1999;159:2461-2468.
- ⁸⁴ Salgado C, Calfee DP, Giannetta ET, Farr BM. Rate of turning culture positive for vancomycin-resistant enterococcus after treatment with oral vancomycin. Presented at the 39th Annual Meeting of the Infectious Diseases Society of America; October 25-28, 2001; San Francisco, CA. Abstract (507):126.
- ⁸⁵ Coque TM, Tomayko JF, Ricke SC, Okhyusen PC, Murray BE. Vancomycin-resistant enterococci from nosocomial, community, and animal sources in the United States. *Antimicrob Agents Chemother* 1996;40:2605-2609.

Appendix B

- ⁸⁶ Morris JG, Shay DK, Hebden JN, et al. Enterococci resistant to multiple antimicrobial agents including vancomycin: establishment of endemicity in a university medical center. *Ann Intern Med* 1995;123:250-259.
- ⁸⁷ Cetinkaya Y, Falk P, Mayhall CG. Vancomycin-resistant enterococci. *Clin Microbiol Rev* 2000;13:686-707.
- ⁸⁸ Bates J. Epidemiology of vancomycin-resistant enterococcus in the community and the relevance of farm animals to human infection. *J Hosp Infect* 1997;37:89-101.
- ⁸⁹ Wegener HC, Aarestrup FM, Jensen LB, Hammerum AM, Bager F. Use of antimicrobial growth promoters in food animals and *Enterococcus faecium* resistance to therapeutic antimicrobial drugs in Europe. *Emerg Infect Dis* 1999;5:329-335.
- ⁹⁰ Klare I, Heier H, Claus H, Reissbrodt R, Witt W. vanA-mediated highlevel glycopeptide resistance in *Enterococcus faecium* from animal husbandry. *FEMS Microbiol Lett* 1995;125:165.
- ⁹¹ Devriese LA, Ieven M, Goossens H, et al. Presence of vancomycin-resistant enterococci in farm and pet animals. *Antimicrob Agents Chemother* 1996;40:2285-2287.
- ⁹² Borgen K, Simonsen G, Sundsfjord A, Wasteson Y, Olsvik O, Kruse H. Continuing high prevalence of VanA-type vancomycin-resistant enterococci on Norwegian poultry farms three years after avoparcin was banned. *J Appl Microbiol* 2000;89:478-485.
- ⁹³ Willems R, Homan W, Top J, et al. Variant esp gene as a marker of a distinct genetic lineage of vancomycin-resistant *Enterococcus faecium* spreading in hospitals. *Lancet* 2001;357:853-855.
- ⁹⁴ Rice LB, Carias L, Rudin S, et al. A potential virulence gene, hylefM, predominates in *Enterococcus faecium* of clinical origin. *J Infect Dis* 2003;187:508-512.
- ⁹⁵ Boyce JM, Mermel LA, Zervos MJ, et al. Controlling vancomycin-resistant enterococci. *Infect Control Hosp Epidemiol* 1995;16:634-637.
- ⁹⁶ Boyce JM, Opal SM, Chow JW, et al. Outbreak of multi-drug resistant *Enterococcus faecium* with transferable vanB class vancomycin resistance. *J Clin Microbiol* 1994;32:1148-1153.
- ⁹⁷ Clark NC, Cooksey RC, Hill BC, Swenson JM, Tenover FC. Characterization of glycopeptide-resistant enterococci from U.S. hospitals. *Antimicrob Agents Chemother* 1993;37:2311-2317.
- ⁹⁸ Handwerker S, Raucher B, Altarac D, et al. Nosocomial outbreak due to *Enterococcus faecium* highly resistant to vancomycin, penicillin, and gentamicin. *Clin Infect Dis* 1993;16:750.
- ⁹⁹ Livornese LL, Dias S, Romanowski B, et al. Hospital-acquired infection with vancomycin-resistant *Enterococcus faecium* transmitted by electronic thermometers. *Ann Intern Med* 1992;117:112-116.
- ¹⁰⁰ Kim WJ, Weinstein RA, Hayden MK. The changing molecular epidemiology and establishment of endemicity of vancomycin resistance in enterococci at one hospital over a 6-year period. *J Infect Dis* 1999;179:163-171.
- ¹⁰¹ Moreno RM, Grota P, Crisp C, et al. Clinical and molecular epidemiology of vancomycin-resistant *Enterococcus faecium* during its emergence in a city in Southern Texas. *Clin Infect Dis* 1995;21:1234-1237.
- ¹⁰² Byers KE, Anglim AM, Anneski CJ, et al. A hospital epidemic of vancomycin-resistant enterococcus: risk factors and control. *Infect Control Hosp Epidemiol* 2001;22:140-147.
- ¹⁰³ Handwerker S, Skoble J, Discotto LF, Pucci MJ. Heterogeneity of the vanA gene in clinical isolates of enterococci from the Northeastern United States. *Antimicrob Agents Chemother* 1995;39:362-368.
- ¹⁰⁴ Mato R, de Lancesstre H, Carraher M, Robers RB, Tomasz A. Multiplicity of genetic backgrounds among vancomycin-resistant *Enterococcus faecium* isolates recovered from an outbreak in a New York City Hospital. *Microb Drug Resist* 1996;2:309-317.
- ¹⁰⁵ Slaughter S, Hayden MK, Nathan C, et al. A comparison of the effect of universal use of gloves and gowns with that of glove use alone on acquisition of vancomycin-resistant enterococci in a medical intensive care unit. *Ann Intern Med* 1996;125:448-456.

Appendix B

- ¹⁰⁶ Muto CA, Posey K, Pokrywka M, et al. The value of identifying the vancomycin resistant enterococci (VRE) reservoir using weekly VRE surveillance culturing (VRESC): “the iceberg melts.” Presented at the 12th Annual Meeting of the Society for Healthcare Epidemiology of America; April 6-9, 2002; Salt Lake City, UT. Abstract.
- ¹⁰⁷ Nelson RR, McGregor KF, Brown AR, Amyes GS, Young H. Isolation and characterization of glycopeptide-resistant enterococci from hospitalized patients over a 30 month period. *J Clin Microbiol* 2000;38:2112- 2116.
- ¹⁰⁸ Stosor V, Kruszynski J, Suriano T, Noskin GA, Peterson LR. Molecular epidemiology of vancomycin-resistant enterococci: a 2-year perspective. *Infect Control Hosp Epidemiol* 1999;20:653-659.
- ¹⁰⁹ de Lencastre H, Brown AE, Chung M, Armstrong D, Tomasz A. Role of transposon Tn5482 in the epidemiology of vancomycin-resistant *Enterococcus faecium* in the pediatric oncology unit of a New York City hospital. *Microb Drug Resist* 1999;5:113-129.
- ¹¹⁰ Beezhold DW, Slaughter S, Hayden MK, et al. Skin colonization with vancomycin-resistant enterococci among hospitalized patients with bacteremia. *Clin Infect Dis* 1997;24:704-706.
- ¹¹¹ Tornieporth NG, Roberts RB, John J, Hafner A, Riley LW. Risk factors associated with vancomycin-resistant *Enterococcus faecium* infection or colonization in 145 matched case patients and control patients. *Clin Infect Dis* 1996;23:767-772.
- ¹¹² Bonten MJ, Hayden MK, Nathan C, et al. Epidemiology of colonization of patients and environment with vancomycin-resistant enterococci. *Lancet* 1996;348:1615-1619.
- ¹¹³ Donskey CJ, Chowdhry T, Hecker M, et al. Effect of antibiotic therapy on the density of vancomycin-resistant enterococci in the stool of colonized patients. *N Engl J Med* 2000;343:1925-1932.
- ¹¹⁴ Bonten MJ, Slaughter S, Ambergen AW, et al. The role of “colonization pressure” in the spread of vancomycin-resistant enterococci: an important infection control variable. *Arch Intern Med* 1998;158:1127- 1132.
- ¹¹⁵ Quale J, Landman D, Saurina G, Atwood E, DiTore V, Patel K. Manipulation of a hospital antimicrobial formulary to control an outbreak of vancomycin-resistant enterococci. *Clin Infect Dis* 1996;23:1020-1025.
- ¹¹⁶ Goetz AM, Rihs JD, Wagener MM, Muder RR. Infection and colonization with vancomycin-resistant *Enterococcus faecium* in an acute care Veterans Affairs Medical Center: a 2-year survey. *Am J Infect Control* 1998;26:558-562.
- ¹¹⁷ Fridkin SK, Steward CD, Edwards JR, et al. Surveillance of antimicrobial use and antimicrobial resistance in United States hospitals: Project ICARE phase 2. *Clin Infect Dis* 1999;29:245-252.
- ¹¹⁸ Lautenbach E, LaRosa LA, Marr AM, Nachamkin I, Bilker WB, Fishman NO. Changes in the prevalence of vancomycin-resistant enterococci in response to antimicrobial formulary interventions: impact of progressive restrictions on use of vancomycin and third-generation cephalosporins. *Clin Infect Dis* 2003;36:440-446.
- ¹¹⁹ Ostrowsky BE, Trick WE, Sohn AH, et al. Control of vancomycin-resistant enterococcus in health care facilities in a region. *N Engl J Med* 2001;344:1427-1433.
- ¹²⁰ Montecalvo MA, Jarvis WR, Uman J, et al. Costs and savings associated with infection control measures that reduced transmission of vancomycin-resistant enterococci in an endemic setting. *Infect Control Hosp Epidemiol* 2001;22:437-442.
- ¹²¹ Mortimer EA, Lipsitz PJ, Wolinsky E, et al. Transmission of staphylococci between newborns. *Am J Dis Child* 1962;104:289-295.
- ¹²² Zachary KC, Bayne PS, Morrison V, Ford DS, Silver LC, Hooper DC. Contamination of gowns, gloves, and stethoscopes with vancomycin-resistant enterococci. *Infect Control Hosp Epidemiol* 2001;22:560-564.
- ¹²³ Cookson B, Peters B, Webster M, Phillips I, Rahman M, Noble W. Staff carriage of epidemic methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 1989;27:1471-1476.
- ¹²⁴ Lacey S, Flaxman D, Scales J, Wilson A. The usefulness of masks in preventing transient carriage of epidemic methicillin-resistant *Staphylococcus aureus* by healthcare workers. *J Hosp Infect* 2001;48:308-311.
- ¹²⁵ Wells CL, Juni BA, Cameron SB, et al. Stool carriage, isolation, and mortality during

Appendix B

- outbreak of vancomycin-resistant enterococci in hospitalized medical and/or surgical patients. *Clin Infect Dis* 1995;21:45-50.
- ¹²⁶ Bonilla HF, Zervos MA, Lyons MJ, et al. Colonization with vancomycin-resistant *Enterococcus faecium*: comparison of a long-term care unit with an acute-care hospital. *Infect Control Hosp Epidemiol* 1997;18:333-339.
- ¹²⁷ Noskin GA, Stosor V, Cooper I, Peterson L. Recovery of vancomycin-resistant enterococci on fingertips and environmental surfaces. *Infect Control Hosp Epidemiol* 1995;16:577-581.
- ¹²⁸ Suh HK, Jeon YH, Song SJ. A molecular epidemiologic study of methicillin-resistant *Staphylococcus aureus* infection in patients undergoing middle ear surgery. *Eur Arch Otorhinolaryngol* 1998;255:347-351.
- ¹²⁹ Opal SM, Mayer KH, Stenberg MJ, et al. Frequent acquisition of multiple strains of methicillin-resistant *Staphylococcus aureus* by healthcare workers in an endemic hospital environment. *Infect Control Hosp Epidemiol* 1990;11:479-485.
- ¹³⁰ Devine J, Cooke RP, Wright EP. Is methicillin-resistant *Staphylococcus aureus* (MRSA) contamination of ward-based computer terminals a surrogate marker for nosocomial MRSA transmission and handwashing compliance. *J Hosp Infect* 2001;48:72-75.
- ¹³¹ Layton MC, Perez M, Heald, Patterson JE. An outbreak of mupirocin-resistant *Staphylococcus aureus* on a dermatology ward associated with an environmental reservoir. *Infect Control Hosp Epidemiol* 1993;14:369-375.
- ¹³² Boyce JM, Potter-Bynoe G, Chenevert C, King T. Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: possible infection control implications. *Infect Control Hosp Epidemiol* 1997;18:622-627.
- ¹³³ Neely AN, Maley MP. Survival of enterococci and staphylococci on hospital fabrics and plastics. *J Clin Microbiol* 2000;38:724-726.
- ¹³⁴ Boyce JM, Chenevert C. Isolation gowns prevent health care workers (HCWs) from contaminating their clothing, and possibly their hands, with methicillin-resistant *Staphylococcus aureus* (MRSA) and resistant enterococci. Presented at the Eighth Annual Meeting of the Society for Healthcare Epidemiology of America; April 5-7, 1998; Orlando, FL. Abstract S74:52.
- ¹³⁵ Puzniak LA, Leet T, Mayfield J, Kollef M, Mundy LM. To gown or not to gown: the effect on acquisition of vancomycin resistant enterococci. *Clin Infect Dis* 2002;35:18-25.
- ¹³⁶ Srinivasan A, Song X, Bower R, et al. A prospective study to determine whether cover gowns in addition to gloves decrease nosocomial transmission of vancomycin-resistant enterococci in an ICU. *Infect Control Hosp Epidemiol* 2002;23:424-428.
- ¹³⁷ Calfee DP, Farr BM. Infection control and cost control in the era of managed care. *Infect Control Hosp Epidemiol* 2002;23:407-410.
- ¹³⁸ Karanfil LV, Murphy M, Josephson A, et al. A cluster of vancomycin-resistant *Enterococcus faecium* in an intensive care unit. *Infect Control Hosp Epidemiol* 1992;13:195-200.
- ¹³⁹ Montecalvo MA, Horowitz H, Gedris C, Carbonaro C, Tenover FC, Issah A. Outbreak of vancomycin-, ampicillin-, and aminoglycosideresistant *Enterococcus faecium* bacteremia in an adult oncology unit. *Antimicrob Agents Chemother* 1994;38:1363-1367.
- ¹⁴⁰ Dembry L, Uzokwe K, Zervos M. Control of endemic glycopeptideresistant enterococci. *Infect Control Hosp Epidemiol* 1996;17:286-292.
- ¹⁴¹ Rupp ME, Marion N, Fey PD, et al. Outbreak of vancomycin-resistant *Enterococcus faecium* in a neonatal intensive care unit. *Infect Control Hosp Epidemiol* 2001;22:301-303.
- ¹⁴² Malik RK, Montecalvo MA, Reale MR, et al. Epidemiology and control of vancomycin-resistant enterococci in a regional neonatal intensive care unit. *Pediatric Infect Dis J* 1999;18:352-356.
- ¹⁴³ Muto CA, Karchmer TB, Cage EG, Durbin LJ, Simonton B, Farr BM. The utility of culturing roommates of patients with vancomycin resistant enterococcus. Presented at the Eighth Annual Meeting of the Society for Healthcare Epidemiology of America; April 5-7, 1998; Orlando, FL. Abstract 76:38.
- ¹⁴⁴ Rubin LG, Tucci V, Cercenado E, Elipoulos G, Isenberg HD. Vancomycin-resistant

Appendix B

- Enterococcus faecium* in hospitalized children. *Infect Control Hosp Epidemiol* 1992;13:700-705.
- ¹⁴⁵ Jochimsen E, Fish L, Manning K, et al. Control of vancomycin-resistant enterococci at a community hospital: efficacy of patient and staff cohorting. *Infect Control Hosp Epidemiol* 1999;20:106-109.
- ¹⁴⁶ Golan Y, Sullivan B, Snyderman DR. Elimination of vancomycin-resistant enterococcus (VRE) transmission in a neonatal intensive care unit (NICU). Presented at the 39th Annual Meeting of the Infectious Diseases Society of America; October 25-28, 2001; San Francisco, CA. Abstract 209:75.
- ¹⁴⁷ Price CS, Paule S, Noskin GA, Peterson LR. Active surveillance reduces vancomycin-resistant enterococci bloodstream isolates. Presented at the 39th Annual Meeting of the Infectious Diseases Society of America; October 25-28, 2001; San Francisco, CA. Abstract 212:75.
- ¹⁴⁸ Siddiqui AH, Harris AD, Hebden J, Wilson PD, Morris JG, Roghmann M. The effect of active surveillance for vancomycin resistant enterococci in high risk units on vancomycin resistant enterococci incidence hospital-wide. *Am J Infect Control* 2002;30:40-43.
- ¹⁴⁹ Calfee DP, Giannetta E, Farr BM. Effective control of VRE colonization using CDC recommendations for detection and isolation. Presented at the 38th Annual Meeting of the Infectious Diseases Society of America; September 7-10, 2000; New Orleans, LA. Abstract 21:44.
- ¹⁵⁰ Ackelsberg J, Kostman J. A laboratory and clinical study of stethoscopes as potential fomites of infection. Presented at the 33rd Annual Meeting of the Infectious Diseases Society of America; September 16- 18, 1995; San Francisco, CA.
- ¹⁵¹ Bernard L, Kereveur A, Durand D, et al. Bacterial contamination of hospital physicians' stethoscopes. *Infect Control Hosp Epidemiol* 1999;20:626-628.
- ¹⁵² Breathnach AS, Jenkins DR, Pedler SJ. Stethoscopes as possible vectors of infection by staphylococci. *BMJ* 1992;305:1573-1574.
- ¹⁵³ Maki DG, Halvorson K, Fisher S. The stethoscope: a medical device with potential for amplifying cross-infection of resistant nosocomial organisms in the hospital. Presented at the 36th Annual Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 15-18, 1996; New Orleans, LA. Abstract 154:247.
- ¹⁵⁴ Smith MA, Mathewson JJ, Ulert IA, Scerpella EG, Ericsson CD. Contaminated stethoscopes revisited. *Arch Intern Med* 1996;156:82-84.
- ¹⁵⁵ Cohen HA, Amir J, Matalon A, Mayan R, Beni S, Barzilai A. Stethoscopes and otoscopes: a potential vector of infection? *Fam Pract* 1997;14:446-449.
- ¹⁵⁶ Singh D, Kaur H, Gardner WG, Treen LB. Bacterial contamination of hospital papers. *Infect Control Hosp Epidemiol* 2002;23:274-276.
- ¹⁵⁷ Brooks S, Khan A, Stoica D, Griffith J. Reduction in vancomycin-resistant *Enterococcus* and *Clostridium difficile* infections following change to tympanic thermometers. *Infect Control Hosp Epidemiol* 1998;19:333-336.
- ¹⁵⁸ Centers for Disease Control and Prevention/Healthcare Infection Control Practices Advisory Committee (HICPAC). *Draft Guidelines for Environmental Infection Control in Healthcare Facilities*. Atlanta, GA: Centers for Disease Control and Prevention; 2001. Available at www.cdc.gov/ncidod/hip/enviro/guide.htm.
- ¹⁵⁹ Edmond MB, Ober JF, Weinbaum DL, et al. Vancomycin-resistant *Enterococcus faecium* bacteremia: risk factors for infection. *Clin Infect Dis* 1995;20:1126-1133.
- ¹⁶⁰ Wendt C, Wiesenthal B, Dietz E, Ruden H. Survival of vancomycin-resistant and vancomycin-susceptible enterococci on dry surfaces. *J Clin Microbiol* 1998;36:3734-3736.
- ¹⁶¹ Smith TL, Iwen PC, Olson SB, Rupp ME. Environmental contamination with vancomycin-resistant enterococci in an outpatient setting. *Infect Control Hosp Epidemiol* 1998;19:515-518.
- ¹⁶² Oie S, Kamiya A. Survival of methicillin-resistant *Staphylococcus aureus* (MRSA) on naturally contaminated dry mop. *J Hosp Infect* 1996;34:145-149.
- ¹⁶³ Noskin GA, Peterson L, Warren J. *Enterococcus faecium* and *Enterococcus faecalis*

Appendix B

- bacteremia: acquisition and outcome. *Clin Infect Dis* 1995;20:296-301.
- ¹⁶⁴ Dietz B, Raht A, Wendt C, Martiny H. Survival of MRSA on sterile goods packaging. *J Hosp Infect* 2001;49:255-261.
- ¹⁶⁵ McDade J, Hall L. Survival of *Staphylococcus aureus* in the environment: I. Exposure on surfaces. *American Journal of Hygiene* 1963;78:330-337.
- ¹⁶⁶ Rutala W, Katz E, Sherertz R, Sarubbi F. Environmental study of methicillin-resistant *Staphylococcus aureus* epidemic in a burn unit. *J Clin Microbiol* 1983;18:683-688.
- ¹⁶⁷ Embil J, McLeod J, Al-Barrak AM, et al. An outbreak of methicillin-resistant *Staphylococcus aureus* on a burn unit: potential role of contaminated hydrotherapy equipment. *Burn* 2001;27:681-688.
- ¹⁶⁸ Noskin GA, Bednarz P, Suriano T, Reiner S, Peterson LR. Persistent contamination of fabric-covered furniture by vancomycin-resistant enterococci: implications for upholstery selection in hospitals. *Am J Infect Control* 2000;28:311-313.
- ¹⁶⁹ Falk PS, Winnike J, Woodmansee C, Desai M, Mayhall CG. Outbreak due to vancomycin-resistant enterococci (VRE) in a burn unit. *Infect Control Hosp Epidemiol* 2000;21:575-582.
- ¹⁷⁰ Rimland D. Nosocomial infections with methicillin and tobramycin resistant *Staphylococcus aureus*: implication of physiotherapy in hospital-wide dissemination. *Am J Med Sci* 1985;290:91-97.
- ¹⁷¹ Law MR, Gill ON. Hospital-acquired infection with methicillin-resistant and methicillin-sensitive staphylococci. *Epidemiol Infect* 1988;101:623-629.
- ¹⁷² Murray-Leisure KA, Geib S, et al. Control of epidemic methicillin-resistant *Staphylococcus aureus*. *Infect Control Hosp Epidemiol* 1990;11:343-350.
- ¹⁷³ Nicolle LE, Dyck B, Thompson G, et al. Regional dissemination and control of epidemic methicillin-resistant *Staphylococcus aureus*. *Infect Control Hosp Epidemiol* 1999;20:202-205.
- ¹⁷⁴ Cantej J, Rhoton B, Southgate W, Snyder C. Control of spread of methicillin resistant *Staphylococcus aureus* in a neonatal ICU. Presented at the 12th Annual Meeting of the Society for Healthcare Epidemiology of America; April 6-9, 2002; Salt Lake City, UT. Abstract 36:49.
- ¹⁷⁵ Croyle K, Muto C. Surveillance cultures in the race against MRSA: ahead by a nose. Presented at the 12th Annual Meeting of the Society for Healthcare Epidemiology of America; April 6-9, 2002; Salt Lake City, UT. Abstract 35:49.
- ¹⁷⁶ Kotilainen P, Routamaa M, Peltonen R, et al. Eradication of methicillin-resistant *Staphylococcus aureus* from a health center ward and associated nursing home. *Arch Intern Med* 2001;161:859-863.
- ¹⁷⁷ Nouer A, Araujo A, Chebabo A, Cardoso F, Pinto M, Hospital Universitário Universidade Federal do Rio de Janeiro. Control of methicillin-resistant *Staphylococcus aureus* (MRSA) in an intensive care unit (ICU) after the institution of routine screening. Presented at the 42nd General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 27-30, 2002; San Francisco, CA. Abstract K-97:97.
- ¹⁷⁸ Horcajada J, Marco F, Martinez J, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* colonization at admission in a tertiary hospital: usefulness of early detection. Presented at the 42nd General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 27-30, 2002; San Francisco, CA. Abstract K-98.
- ¹⁷⁹ Gerard M, Dediste A, Van Esse R, et al. Cost effectiveness of a policy of methicillin resistant *Staphylococcus aureus* (MRSA) screening, decontamination and isolation in a medical ICU. Presented at the 42nd General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 27-30, 2002; San Francisco, CA. Abstract K-99.
- ¹⁸⁰ Green K, Fleming CA, Richardson H, Low DE, Willey B, McGeer A. MRSA and VRE in Ontario, Canada: results of 7 years of surveillance and control measures. Presented at the 42nd General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 27-30, 2002; San Francisco, CA. Abstract K661.
- ¹⁸¹ Muto CA, Giannetta ET, Durbin LJ, Simonton BM, Farr BM. Cost effectiveness of peri-

Appendix B

- rectal surveillance cultures for controlling vancomycin-resistant enterococcus. *Infect Control Hosp Epidemiol* 2002;23:429-435.
- ¹⁸² Armstrong-Evans M, Litt M, Willey B, et al. Control of transmission of vancomycin-resistant *Enterococcus faecium* in a long-term-care facility. *Infect Control Hosp Epidemiol* 1999;20:312-317.
- ¹⁸³ Hill AB. *A Short Textbook of Medical Statistics*, vol. 11. London: Hodder and Stoughton; 1984.
- ¹⁸⁴ Salgado C, Sherertz R, Karchmer T, et al. Public health initiative to control MRSA and VRE in Virginia and North Carolina. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 164:75.
- ¹⁸⁵ Arnold MS, Dempsey JM, Fishman M, McAuley PJ, Tibert C, Vallande NC. The best hospital practices for controlling methicillin-resistant *Staphylococcus aureus*: on the culturing edge. *Infect Control Hosp Epidemiol* 2002;23:69-76.
- ¹⁸⁶ D'Agata EMC, Thayer V, Shaffner W. An outbreak of *Acinetobacter baumannii*: the importance of cross transmission. *Infect Control Hosp Epidemiol* 2000;21:588-591.
- ¹⁸⁷ Simor AE, Lee M, Vearncombe M, et al. An outbreak due to multiresistant *Acinetobacter baumannii* in an acute burn care unit: risk factors for acquisition and management. *Infect Control Hosp Epidemiol* 2002;23:261-267.
- ¹⁸⁸ Piagnerelli M, Carlier E, Deplano A, Lejeune P, Govaerts D. Risk factors for infection and molecular typing in patients in the intensive care unit colonized with nosocomial *Enterobacter aerogenes*. *Infect Control Hosp Epidemiol* 2002;23:452-456.
- ¹⁸⁹ Selkon JB, Stokes ER, Ingham HR. The role of an isolation unit in the control of hospital infection with methicillin resistant staphylococci. *J Hosp Infect* 1980;1:41-46.
- ¹⁹⁰ Pearman JW, Christiansen KJ, Annear DI, et al. Control of methicillin resistant *Staphylococcus aureus* (MRSA) in an Australian metropolitan Vol. 24 No. 5 PREVENTING SPREAD OF ANTIBIOTIC RESISTANCE 383 teaching hospital complex. *Med J Aust* 1985;142:103-108.
- ¹⁹¹ Shanson DC, Johnstone D, Midgley J. Control of a hospital outbreak of methicillin-resistant *Staphylococcus aureus* infections: value of an isolation unit. *J Hosp Infect* 1985;6:285-292.
- ¹⁹² Price EH, Brain A, Dickson JAS. An outbreak with a gentamicin and methicillin-resistant *Staphylococcus aureus* in a neonatal unit. *J Hosp Infect* 1980;1:221-228.
- ¹⁹³ Farrington M, Redpath C, Trundle C, Coomber S, Brown NM. Winning the battle but losing the war: methicillin-resistant *Staphylococcus aureus* (MRSA) infection at a teaching hospital. *QJM* 1998;91:539-548.
- ¹⁹⁴ Kunin CM, Tupasi T, Craig WA. Use of antibiotics: a brief exposition of the problem and some tentative solutions. *Ann Intern Med* 1973;79:555-560.
- ¹⁹⁵ Zuckerman RA, Steele L, Venezia RA, Tobin EH. Undetected vancomycin-resistant *Enterococcus* in surgical intensive care unit patients. *Infect Control Hosp Epidemiol* 1999;20:685-686.
- ¹⁹⁶ Ostrowsky B, Venkataraman L, D'Agata E, Gold H, DeGirolami P, Samore M. Vancomycin-resistant enterococci in intensive care units: high frequency of stool carriage during a non-outbreak period. *Arch Intern Med* 1999;159:1467-1472.
- ¹⁹⁷ Austin DJ, Bonten MJM, Weinstein RA, Slaughter S, Anderson RM. Vancomycin-resistant enterococci in intensive-care hospital settings: transmission dynamics, persistence, and the impact of infection control programs. *Proc Natl Acad Sci U S A* 1999;96:6908-6913.
- ¹⁹⁸ Sebile V, Chevret S, Valleron A. Modeling the spread of resistant nosocomial pathogens in an intensive care unit. *Infect Control Hosp Epidemiol* 1997;18:84-92.
- ¹⁹⁹ Warren DK, Kollef MH, Seiler SM, Fridken SK, Fraser VJ. The epidemiology of vancomycin-resistant *Enterococcus* colonization in a medical intensive care unit. *Infect Control Hosp Epidemiol* 2003;24:257-263.
- ²⁰⁰ Ridwan B, Mascini E, van Der RN, Verhoef J, Bonten M. What action should be taken to prevent spread of vancomycin resistant enterococci in European hospitals? *BMJ* 2002;324:666-668.

Appendix B

- ²⁰¹ Brisse S, Fussing V, Ridwan B, Verhoef J, Willems RJ. Automated ribotyping of vancomycin resistant *Enterococcus faecium* isolates. *J Clin Microbiol* 2002;40:1977-1984.
- ²⁰² Francois P, Pittet D, Bento M, et al. Rapid detection of methicillin-resistant *Staphylococcus aureus* directly from sterile or nonsterile clinical samples by a new molecular assay. *J Clin Microbiol* 2003;41:254-260.
- ²⁰³ Lankford MG, Zembower TR, Trick WE, Hacek DM, Noskin GA, Peterson LR. Impact of hospital design on the handwashing compliance among healthcare workers. Presented at the 38th Annual Meeting of the Infectious Diseases Society of America; September 7- 10, 2000; New Orleans, LA. Abstract 18:43.
- ²⁰⁴ Preston GA, Larson EL, Stamm W. The effect of private isolation rooms on patient care practices, colonization and infection in an intensive care unit. *Am J Med* 1981;70:641-645.
- ²⁰⁵ Albert RK, Condie F. Handwashing patterns in medical intensive care units. *N Engl J Med* 1981;304:1465.
- ²⁰⁶ Larson E. Compliance with isolation technique. *Am J Infect Control* 1983;11:221-225.
- ²⁰⁷ Donowitz LG, Hunt EH, Pugh VG, Farr BM, Hendley JO. Comparison of historical and serologic immunity to varicella-zoster virus in 373 hospital employees. *Am J Infect Control* 1987;15:212-214.
- ²⁰⁸ Graham M. Frequency and duration of handwashing in an intensive care unit. *Am J Infect Control* 1990;18:77-81.
- ²⁰⁹ Dubbert PM, Dolce J, Richter W, Miller M, Chapman S. Increasing ICU staff handwashing: effects of education and group feedback. *Infect Control Hosp Epidemiol* 1990;11:191-193.
- ²¹⁰ Pettinger A, Nettleman MD. Epidemiology of isolation precautions. *Infect Control Hosp Epidemiol* 1991;12:303-307.
- ²¹¹ Larson EL, McGinley K, Foglia A, et al. Handwashing practices and resistance and density of bacterial hand flora on two pediatric units in Lima, Peru. *Infect Control Hosp Epidemiol* 1992;20:65-72.
- ²¹² Doebbeling BN, Pfaller MA, Houston AK, Wenzel RP. Removal of nosocomial pathogens from the contaminated glove: implications for glove reuse and handwashing. *Ann Intern Med* 1988; 109:394-398.
- ²¹³ Zimakoff J, Kjelsberg AB, Larsen SO, Holstein B. A multicenter questionnaire investigation of attitudes toward hand hygiene, assessed by the staff in fifteen hospitals in Denmark and Norway. *Am J Infect Control* 1992;20:58-64.
- ²¹⁴ Meengs MR, Giles BK, Chisholm CD, Cordell WH, Nelson DR. Hand washing frequency in an emergency department. *J Emerg Nurs* 1994;20:183-188.
- ²¹⁵ Pittet D, Hugonnet S, Harbarth S, et al. Effectiveness of a hospitalwide programme to improve compliance with hand hygiene: infection control programme. *Lancet* 2000;356:1307-1312.
- ²¹⁶ Pittet D, Mourouga P, Perneger TV. Compliance with handwashing in a teaching hospital. *Ann Intern Med* 1999;130:126-130.
- ²¹⁷ Vernon MO, Trick WE, Welbel SF, Peterson BJ, Weinstein RA. Hand hygiene adherence: does the number of sinks matter? *Infect Control Hosp Epidemiol* 2003;24:224-225.
- ²¹⁸ Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings: recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *MMWR* 2002;51(RR16):1-45.
- ²¹⁹ Grundmann H, Hori S, Winter B, Tami A, Austin DJ. Risk factors for the transmission of methicillin-resistant *Staphylococcus aureus* in an adult intensive care unit: fitting a model to the data. *J Infect Dis* 2002;185:481-488.
- ²²⁰ Larson EL, Early E, Cloonan P, Sugrue S, Parides M. An organizational climate intervention associated with increased handwashing and decreased nosocomial infections. *Behav Med* 2000;26:14-22.
- ²²¹ Webster J, Faoagali JL, Cartwright D. Elimination of methicillin-resistant *Staphylococcus aureus* from a neonatal intensive care unit after hand washing with triclosan. *J Paediatr Child Health* 1994;30:59-64.
- ²²² Zafar AB, Butler RC, Reese DJ, Gaydos LA. Use of 0.3% triclosan (Bacti-Stat®) to eradi-

Appendix B

- cate an outbreak of methicillin-resistant *Staphylococcus aureus* in a neonatal nursery. *Am J Infect Control* 1995;23:200-208.
- ²²³ Lucet JC, Rigaud MP, Mentre F, et al. Hand contamination before and after different hand hygiene techniques: a randomized clinical trial. *J Hosp Infect* 2002;50:276-280.
- ²²⁴ Pittet D, Dharan S, Touveneau S, Sauvan V, Perneger TV. Bacterial contamination of the hands of hospital staff during routine patient care. *Arch Intern Med* 1999;159:821-826.
- ²²⁵ Klein BS, Perloff WH, Maki DG. Reduction of nosocomial infection during pediatric intensive care by protective isolation. *N Engl J Med* 1989;320:1714-1721.
- ²²⁶ Leclair JM, Freeman J, Sullivan BF, Crowley CM, Goldmann DA. Prevention of nosocomial RSV infections through compliance with glove and gown isolation precautions. *N Engl J Med* 1987;317:329-334.
- ²²⁷ Olsen RJ, Lynch P, Coyle MB, Cummings J, Bokete T, Stamm WE. Examination gloves as barriers to hand contamination in clinical practice. *JAMA* 1993;270:350-353.
- ²²⁸ Tenorio AR, Badri SM, Sahgal NB, et al. Effectiveness of gloves in the prevention of hand carriage of vancomycin-resistant enterococcus species by health care workers after patient care. *Clin Infect Dis* 2001;32:826-829.
- ²²⁹ Johnson S, Gerding DN, Olson MM, et al. Prospective, controlled study of vinyl glove use to interrupt *Clostridium difficile* nosocomial transmission. *Am J Med* 1990;88:137-140.
- ²³⁰ Muto CA, Byers KE, Karchmer TB, Dill JB, Durbin LJ, Farr BM. Controlling vancomycin-resistant enterococcus (VRE) at a university hospital. Presented at the Seventh Annual Meeting of the Society for Healthcare Epidemiology of America; April 27-29, 1997; St. Louis, MO. Abstract 52:28.
- ²³¹ Karchmer TB, Ribadeneyra MG, Durbin LJ, Giannetta E, Farr BM. Prevalence of methicillin-resistant *Staphylococcus aureus* colonization among resident-physicians at a teaching hospital using contact/droplet precautions for MRSA isolation. Presented at the Tenth Annual Meeting of the Society for Healthcare and Epidemiology of America; March 5-9, 2000; Atlanta, GA. Abstract 200.
- ²³² Sherertz RJ, Reagan DR, Hampton KD, et al. A cloud adult: the *Staphylococcus aureus*-virus interaction revisited. *Ann Intern Med* 1996;124:539-547.
- ²³³ Williams RE. Epidemiology of airborne staphylococcal infection. *Bacteriol Rev* 1966;30:660-674.
- ²³⁴ Mermel L, Dempsey J, Parenteau S. An MRSA outbreak in a surgical intensive care unit: possible role of aerosol transmission from opened ventilator tubing. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 17:42.
- ²³⁵ Meester M, Schultz C, Boeijen-Donkers L, et al. Evidence for airborne transmission of methicillin-resistant *S. aureus*. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 36:46.
- ²³⁶ Bischoff W, Bassetti S, Bassetti-Wyss B, et al. 'The cloud phenomenon' (CP): predictors of *Staphylococcus aureus* (SA) airborne dispersal associated with rhinovirus infection. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 100:60.
- ²³⁷ Shiomori T, Miyamoto H, Makishima K. Significance of airborne transmission of methicillin-resistant *Staphylococcus aureus* in an otolaryngology- head and neck surgery unit. *Arch Otolaryngol Head Neck Surg* 2001;127:644-648.
- ²³⁸ Anonymous. *Staphylococcus aureus* resistant to vancomycin: United States, 2002. *MMWR* 2002;51:565-567. 384 INFECTION CONTROL AND HOSPITAL EPIDEMIOLOGY May 2003
- ²³⁹ Achong M, Hauser BA, Krusky JL. Rational and irrational use of antibiotics in a Canadian teaching hospital. *Canadian Medical Association Journal* 1977;116:256.
- ²⁴⁰ Kunin CM. The responsibility of the infectious disease community for the optimal use of antimicrobial agents. *J Infect Dis* 1985;151:388.
- ²⁴¹ Maki DG, Schuna AA. A study of antimicrobial misuse in a university hospital. *Am J Med Sci* 1978;275:271.

Appendix B

- ²⁴² Roberts AW, Visconti JA. The rational and irrational use of systemic antimicrobial drugs. *American Journal of Hospital Pharmacy* 1972;29:828.
- ²⁴³ Scheckler WE, Bennett JV. Antibiotic use in seven community hospitals. *JAMA* 1970;214:264.
- ²⁴⁴ Bamberger DM, Dahl SL. Impact of voluntary vs enforced compliance of third generation cephalosporin use in a teaching hospital. *Arch Intern Med* 1992;152:554-557.
- ²⁴⁵ Shah SS, Sinkowitz-Cochran, Keyserling H, Jarvis WR. Vancomycin use in pediatric cardiothoracic surgery patients. *Pediatr Infect Dis J* 1999;18:558-560.
- ²⁴⁶ Shah SS, Sinkowitz-Cochran, Keyserling H, Jarvis WR. Vancomycin use in pediatric neurosurgery patients. *Am J Infect Control* 1999;27:482-497.
- ²⁴⁷ Lawton RM, Fridkin SK, Gaynes RP, McGowan JE. Practices to improve antimicrobial use at 47 hospitals: the status of the 1997 SHEA/IDSA position paper recommendations. *Infect Control Hosp Epidemiol* 2000;21:256-259.
- ²⁴⁸ Hopkins HA, Sinkowitz-Cochran RL, Rudin BA, Keyserling HL, Jarvis WR. Vancomycin use in pediatric hematology-oncology patients. *Infect Control Hosp Epidemiol* 2000;21:48-50.
- ²⁴⁹ Shales DM, Gerding DN, John JF, et al. Society for Healthcare Epidemiology of America and Infectious Diseases Society of America Joint Committee on the Prevention of Antimicrobial Resistance: guidelines for the prevention of antimicrobial resistance in hospitals. *Clin Infect Dis* 1997;25:584-599.
- ²⁵⁰ Forrest A, Nix DE, Ballow CH, Goss TF, Birmingham MC, Schentag JJ. Pharmacodynamics of intravenous ciprofloxacin in seriously ill patients. *Antimicrob Agents Chemother* 1993;37:1073-1081.
- ²⁵¹ Monnet DL. Methicillin-resistant *Staphylococcus aureus* and its relationship to antimicrobial use: possible implications for control. *Infect Control Hosp Epidemiol* 1998;19:552-559.
- ²⁵² Weber SG, Gold HS, Karchmer AW, Hooper DC, Carmeli Y. Exposure to quinolones is a risk factor for methicillin-resistant (MRSA), but not for methicillin-susceptible *Staphylococcus aureus* (MSSA). Presented at the 42nd Annual General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 27-30, 2002; San Francisco, CA. Abstract K-2070.
- ²⁵³ Dziekan G, Hahn A, Thune K, et al. Methicillin-resistant *Staphylococcus aureus* in a teaching hospital: investigation of nosocomial transmission using a matched case-control study. *J Hosp Infect* 2000;46:263-270.
- ²⁵⁴ Harbarth S, Harris AD, Carmeli Y, Samore MH. Parallel analysis of individual and aggregated data on antibiotic exposure and resistance in gram-negative bacilli. *Clin Infect Dis* 2001;33:1462-1468.
- ²⁵⁵ Hori S, Sunley R, Tami A, Grundmann H. The Nottingham *Staphylococcus aureus* population study: prevalence of MRSA among elderly in a university hospital. *J Hosp Infect* 2002;50:25-19.
- ²⁵⁶ Campillo B, Dupeyron C, Richardet JP. Epidemiology of hospital-acquired infections in cirrhotic patients: effect of carriage of methicillin-resistant *Staphylococcus aureus* and influence of previous antibiotic therapy and norfloxacin prophylaxis. *Epidemiol Infect* 2001;127:443-450.
- ²⁵⁷ Gruson D, Hilbert G, Vargas F, et al. Rotation and restricted use of antibiotics in a medical intensive care unit: impact on the incidence of ventilator-associated pneumonia caused by antibiotic-resistant gram-negative bacteria. *Am J Respir Crit Care Med* 2000;162:837-843.
- ²⁵⁸ Landman D, Chockalingam M, Quale JM. Reduction in the incidence of methicillin-resistant *Staphylococcus aureus* and ceftazidime-resistant *Klebsiella pneumoniae* following changes in a hospital antibiotic formulary. *Clin Infect Dis* 1999;28:1062-1066.
- ²⁵⁹ Van der Auwera P, Pensart N, Korten V, Murray BE, Leclercq R. Influence of oral glycopeptide on the fecal flora of human volunteers: selection of highly glycopeptide-resistant enterococci. *J Infect Dis* 1996;173:1129-1136.
- ²⁶⁰ Tucci V, Haran MA, Isenberg H. Epidemiology and control of vancomycin-resistant

Appendix B

- enterococci in an adult and children's hospital. *Am J Infect Control* 1997;25:371-376.
- ²⁶¹ Uttley A, Collins C, Naidoo J, George R. Vancomycin-resistant enterococci. *Lancet* 1988;1:57-58.
- ²⁶² Anglim AM, Klym B, Byers KE, Farr BM. Effect of a vancomycin restriction policy on ordering practices during an outbreak of vancomycin-resistant *Enterococcus faecium*. *Arch Intern Med* 1997;157:1132-1136.
- ²⁶³ Shay DK, Maloney SA, Montecalvo M, et al. Epidemiology and mortality risk of vancomycin-resistant enterococcal bloodstream infections. *J Infect Dis* 1995;172: 993-1000.
- ²⁶⁴ Weinstein JW, Roe M, Towns M, et al. Resistant enterococci: a prospective study of prevalence, incidence, and factors associated with colonization in a university hospital. *Infect Control Hosp Epidemiol* 1996;17:36-41.
- ²⁶⁵ Carmeli Y, Eliopoulos GM, Samore MH. Antecedent treatment with different antibiotics as a risk for vancomycin resistant enterococcus. *Emerg Infect Dis* 2002;8:802-807.
- ²⁶⁶ Lucas GM, Lechtzin N, Puryear DW, Yau LL, Moore RD. Vancomycin-resistant and vancomycin-susceptible enterococcal bacteremia: comparison of clinical features and outcomes. *Clin Infect Dis* 1998;26:1127-1133.
- ²⁶⁷ Donskey CJ, Hanrahan JA, Hutton RA, Rice LB. Effect of parenteral antibiotic administration on the establishment of colonization with vancomycin-resistant *Enterococcus faecium* in the mouse gastrointestinal tract. *J Infect Dis* 2000;181:1830-1833.
- ²⁶⁸ Bradley SJ, Wilson ALT, Allen MC, Sher HA, Goldstone AH, Scott GM. The control of hyperendemic glycopeptide-resistant *Enterococcus* spp. on a haematology unit by changing antibiotic usage. *J Antimicrob Chemother* 1999;43:261-266.
- ²⁶⁹ Fridkin SK, Lawton R, Edwards JR, Tenover FC, McGowan JE, Gaynes RP. Monitoring antimicrobial use and resistance: comparison with a national benchmark on reducing vancomycin use and vancomycin-resistant enterococci. *Emerg Infect Dis* 2002;8:7.
- ²⁷⁰ Donskey CJ, Hanrahan JA, Hutton RA, Rice LB. Effect of parenteral antibiotic administration on persistence of vancomycin-resistant *Enterococcus faecium* in the mouse gastrointestinal tract. *J Infect Dis* 1999;180:384-390.
- ²⁷¹ Struelens MJ, Ronveaux O, Jans B, Mertens R, the Groupement pour le Depistage, 'Etude et la Prevention des Infections Hospitalieres. Methicillin-resistant *Staphylococcus aureus* epidemiology and control in Belgian hospitals, 1991 to 1995. *Infect Control Hosp Epidemiol* 1996;17:503-508.
- ²⁷² Karchmer TB, Jernigan JA, Durbin BM, Simonton BM, Farr BM. Eradication of methicillin-resistant *S. aureus* (MRSA) colonization with different regimens. Presented at the Ninth Annual Meeting of the Society for Healthcare Epidemiology of America; April 18-20, 1999; San Francisco, CA. Abstract 65:42.
- ²⁷³ Bartzokas CA, Paton JH, Gibson MF, Graham R, McLoughlin GA, Croton RS. Control and eradication of methicillin-resistant *Staphylococcus aureus* on a surgical unit. *N Engl J Med* 1984;10:255-259.
- ²⁷⁴ Tuffnell DJ, Croton RS, Hemingway DM, Hartley MN, Wake PN, Garvey RJP. Methicillin-resistant *Staphylococcus aureus*: the role of antisepsis in the control of an outbreak. *J Hosp Infect* 1987;10:255-259.
- ²⁷⁵ Boyce JM, Opal SM, Potter-Bynoe G, Mederios AA. Spread of methicillin-resistant *Staphylococcus aureus* in a hospital and after exposure to a health-care worker with chronic sinusitis. *Clin Infect Dis* 1993;17:496-504.
- ²⁷⁶ Kluytmans J, van Leeuwen W, Goessens W, et al. Food-initiated outbreak of methicillin-resistant *Staphylococcus aureus* analyzed by phenoand genotyping. *J Clin Microbiol* 1995;33:1121-1128.
- ²⁷⁷ Lessing MP, Jordens JZ, Bowlwer IC. When should health care workers be screened for methicillin resistant *Staphylococcus aureus*? *J Hosp Infect* 1997;35:320-321.
- ²⁷⁸ Doebbeling BN, Breneman DL, Neu HC, et al. Elimination of *Staphylococcus aureus* nasal carriage in health care workers: analysis of six clinical trials with mupirocin calcium ointment. The Mupirocin Collaborative Study Group. *Clin Infect Dis* 1993;17:466-474.
- ²⁷⁹ Doebbeling BN, Regan DR, Pfaller MA, Houston AK, Hollis RJ, Wenzel RP. Long term

Appendix B

- efficacy of intranasal mupirocin ointment: a prospective cohort study of *Staphylococcus aureus* carriage. *Arch Intern Med* 1994;154:1505-1508.
- ²⁸⁰ Regan DR, Doebbeling BN, Pfaller MA, et al. Elimination of coincident *Staphylococcus aureus* nasal and hand carriage with intranasal application of mupirocin ointment: a prospective cohort study of *Staphylococcus aureus* carriage. *Ann Intern Med* 1991;114:101-106.
- ²⁸¹ Perl TM, Cullen JJ, Wenzel RP, et al. Intranasal mupirocin to prevent postoperative *Staphylococcus aureus* infections. *N Engl J Med* 2002;13:1871-1877.
- ²⁸² Farr BM. Mupirocin to prevent *S. aureus* infections. *N Engl J Med* 2002;13:1905-1906.
- ²⁸³ Harbarth S, Dharan S, Liassine N, Herrault P, Auckenthaler R, Pittet D. Randomized, placebo-controlled, double-blind trial to evaluate the efficacy of mupirocin for eradicating carriage of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1999;43:1412-1416.
- ²⁸⁴ Harbarth S, Dharan S, Liassine N, Herrault P, Auckenthaler R, Pittet D. 24 No. 5 PREVENTING SPREAD OF ANTIBIOTIC RESISTANCE 385 D. Risk factors for persistent carriage of methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis* 2000;31:1380-1385.
- ²⁸⁵ Vasquez JE, Walker ES, Franzus BW, Overbay BK, Regan DR, Sarubbi FA. The epidemiology of mupirocin resistance among methicillin-resistant *Staphylococcus aureus* at a Veterans' Affairs hospital. *Infect Control Hosp Epidemiol* 2000;21:459-464.
- ²⁸⁶ Miller MA, Dascal A, Portnory J, Medelson J. Development of mupirocin resistance among methicillin-resistant *Staphylococcus aureus* after widespread use of nasal mupirocin ointment. *Infect Control Hosp Epidemiol* 1996;17:811-813.
- ²⁸⁷ Centers for Disease Control and Prevention. Recommendations for preventing the spread of vancomycin resistance: recommendations of the Hospital Infection Control Practices Advisory Committee (HICPAC). *MMWR* 1995;44:1-13.
- ²⁸⁸ Centers for Disease Control and Prevention. Guidelines for handwashing and hospital environmental control. *MMWR* 1998;36(suppl):2S.
- ²⁸⁹ Spaulding EH. Chemical disinfection of medical and surgical materials. In: Lawrence CA, Block SS, eds. *Disinfection, Sterilization, and Preservation*. Philadelphia: Lea & Febiger; 1968:517-531.
- ²⁹⁰ Rutala W. APIC guidelines for selection and use of disinfectants. *Am J Infect Control* 1996;24:313-342.
- ²⁹¹ Saurina G, Landman D, Quale J. Activity of disinfectants against vancomycin-resistant *Enterococcus faecium*. *Infect Control Hosp Epidemiol* 1998;18:345-347.
- ²⁹² Rutala W, Stiegel M, Sarubbi F, Weber D. Susceptibility of antibiotic-susceptible and antibiotic-resistant hospital bacteria to disinfectants. *Infect Control Hosp Epidemiol* 1997;18:417-421.
- ²⁹³ Anderson RL, Carr JH, Bond WW, Favero MS. Susceptibility of vancomycin-resistant enterococci to environmental disinfectants. *Infect Control Hosp Epidemiol* 1997;18:195-199.
- ²⁹⁴ Byers KE, Durbin LJ, Simonton BM, Anglim AM, Adal KA, Farr BM. Disinfection of hospital rooms contaminated with vancomycin-resistant *Enterococcus faecium*. *Infect Control Hosp Epidemiol* 1998;19:261-264.
- ²⁹⁵ Rampling A, Wiseman S, Davis L, et al. Evidence that hospital hygiene is important in the control of methicillin resistant *Staphylococcus aureus*. *J Hosp Infect* 2001;49:109-116.
- ²⁹⁶ Mangi RJ, Andriole VT. Contaminated stethoscopes: a potential source of nosocomial infections. *Yale J Biol Med* 1972;45:600-604.
- ²⁹⁷ Pittet D, Safran E, Harbarth S, et al. Automatic alerts for methicillin-resistant *Staphylococcus aureus* surveillance and control: role of a hospital information system. *Infect Control Hosp Epidemiol* 1996;17:496-502.
- ²⁹⁸ Wakefield DS, Helms CM, Massanari RM, Mori M, Pfaller MA. Cost of nosocomial infection: relative contributions of laboratory, antibiotic, and per diem costs in serious *Staphylococcus aureus* infections. *Am J Infect Control* 1988;16:185-192.
- ²⁹⁹ Wakefield DA, Pfaller MA, Hammons GT, Massanari RM. Use of the appropriateness

Appendix B

- evaluation protocol for estimating the incremental costs associated with nosocomial infections. *Med Care* 1987;25:481- 488.
- ³⁰⁰ Arnow PM, Quimosing EM, Beach M. Consequences of intravascular sepsis. *Clin Infect Dis* 1993;16:778-784.
- ³⁰¹ Rubin RJ, Harrington CA, Poon A, Dietrich K, Greene JA, Moiduddin A. The economic impact of *Staphylococcus aureus* infection in New York City hospitals. *Emerg Infect Dis* 1999;5:9-17.
- ³⁰² Abramson MA, Sexton DJ. Nosocomial methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* primary bacteremia: at what costs? *Infect Control Hosp Epidemiol* 1999;20:408-411.
- ³⁰³ Cheng AF, French GL. Methicillin-resistant *Staphylococcus aureus* bacteremia in Hong Kong. *J Hosp Infect* 1988;12:91-101.
- ³⁰⁴ Stone PW, Larson E, Kawar LN. A systematic audit or economic evidence linking nosocomial infections and infection control interventions: 1990-2000. *Am J Infect Control* 2002;30:145-152.
- ³⁰⁵ Kaye KS, Engemann JJ, Mozaffari E, Carmeli Y. Outcomes related to vancomycin resistant *Enterococcus* and methicillin resistant *Staphylococcus aureus*: a comparison of the two types of control groups. Presented at the 42nd Annual General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 27-30, 2002; San Francisco, CA.
- ³⁰⁶ Stosor V, Peterson L, Postelnick M, Noskin G. *Enterococcus faecium* bacteremia: does vancomycin resistance make a difference? *Arch Intern Med* 1998;158:522-527.
- ³⁰⁷ Jernigan JA, Hadziyannis SC. Vancomycin-resistant *Enterococcus faecium* (VRE) bacteremia (B) in severely neutropenic patients. Presented at the 36th Annual General Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy; September 15-18, 1996; New Orleans, LA. Abstract J8:219.
- ³⁰⁸ Edmond MB, Jones RN, Pfaller MA, Wallace SE, Wenzel RP. Multicenter surveillance for nosocomial enterococcal bacteremia: a comparison of vancomycin-sensitive vs vancomycin-resistant cases. Presented at the Sixth Annual Meeting of the Society for Healthcare Epidemiology of America; April 21-23, 1996; Washington, DC. Abstract 17:18.
- ³⁰⁹ Lautenbach E, Bilker WB, Brennan PJ. Enterococcal bacteremia: risk factors for vancomycin resistance and predictors of mortality. *Infect Control Hosp Epidemiol* 1999;20:318-323.
- ³¹⁰ Stroud L, Edwards J, Danzing L, Culver D, Gaynes R. Risk factors for mortality associated with enterococcal bloodstream infections. *Infect Control Hosp Epidemiol* 1996;17:576-580.
- ³¹¹ Newel KA, Millis JM, Arnow PM, et al. Incidence and outcome of infection by vancomycin-resistant enterococcus following orthotopic liver transplantation. *Transplant* 1998;65:439-442.
- ³¹² Bhavnani SM, Drake JA, Forrest A, et al. A nationwide, multicenter, case-control study comparing risk factors, treatment, and outcome for vancomycin-resistant and -susceptible enterococcal bacteremia. *Diagn Microbiol Infect Dis* 2000;36:145-158.
- ³¹³ Song X, Srinivisan A, Plaut D, Perl TM. Effect of nosocomial vancomycin-resistant enterococcal bacteremia on mortality, length of stay, and cost. *Infect Control Hosp Epidemiol* 2003;24:251-256.
- ³¹⁴ Karchmer TB, Cage EG, Durbin LJ, Simonton BM, Farr BM. Cost effectiveness of active surveillance cultures for controlling methicillin-resistant *S. aureus* (MRSA). *J Hosp Infect* 2002;51:126-132.
- ³¹⁵ Papia G, Louie M, Tralla A, Johnson C, Collins V, Simor AE. Screening high-risk patients for methicillin-resistant *Staphylococcus aureus* on admission to the hospital: is it cost effective? *Infect Control Hosp Epidemiol* 1999;20:473-477.
- ³¹⁶ Bronstein M, Kaye K, Sexton D. Gown utilization as a measure of cost of methicillin resistant *Staphylococcus aureus* screening. Presented at the 12th Annual Meeting of the Society for Healthcare Epidemiology of America; April 6-9, 2002; Salt Lake City, UT. Abstract: 47:51.

Appendix B

- ³¹⁷ Lucet J, Chevret S, Durand-Zaleski I, Chastang Cregnier B. Prevalence and risk factors for carriage of methicillin resistant *Staphylococcus aureus* at admission to the intensive care unit. *Arch Intern Med* 2003;163:181-188.
- ³¹⁸ Karchmer TB, Farr BM. Presumptive isolation on admission of patients transferred from facilities with a high-prevalence of MRSA: a cost analysis. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 38:46.
- ³¹⁹ Fridken SK, Hageman J, McDougal LK, et al. Epidemiological and microbiological characterization of infections caused by *Staphylococcus aureus* with reduced susceptibility to vancomycin: United States 1997- 2001. *Clin Infect Dis* 2003;36:429-439.
- ³²⁰ Calfee DP, Giannetta E, Durbin LJ, Farr BM. The increasing prevalence of MRSA and VRE colonization among patients transferred from primary and secondary health care facilities. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 171.
- ³²¹ Muto CA, Cage E, Durbin LJ, Simonton BM, Farr BM. The utility of culturing patients on admission transferred from other hospitals or nursing homes for vancomycin resistant *Enterococcus* (VRE). Presented at the 35th Annual Meeting of the Infectious Diseases Society of America; November 12-15, 1998; Denver, CO. Abstract.
- ³²² Muto CA, Patel-Brown S, Krystofiak S, et al. Vancomycin-resistant enterococci (VRE): hospital-wide point prevalence study and epidemiologic description. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract.
- ³²³ Silverblatt FJ, Tibert C, Mikolich D, et al. Preventing the spread of vancomycin-resistant enterococci in a long-term care facility. *J Am Geriatr Soc* 2000;48:1211-1215.
- ³²⁴ Salgado C, Sheppe S, Dill J, Durbin L, Farr B. The sensitivity of MRSA follow-up cultures. Presented at the 11th Annual Meeting of the Society for Healthcare Epidemiology of America; April 1-3, 2001; Toronto, Ontario, Canada. Abstract 40.
- ³²⁵ Manian FA, Finkle D, Zack J, Meyer L. Routine screening for methicillin- resistant *Staphylococcus aureus* among patients newly admitted to an acute rehabilitation unit. *Infect Control Hosp Epidemiol* 2002;23:516- 519.
- ³²⁶ Larson E, Killien M. Factors influencing handwashing behavior of patient care personnel. *Am J Infect Control* 1982;10:93-99.
- ³²⁷ Maki DG. The use of antiseptics for handwashing by medical personnel. *J Chemother* 1989;1(suppl):3-11.
- ³²⁸ Massanari RM, Hierholzer W Jr. A crossover comparison of antiseptic soaps on nosocomial infection rates in intensive care units. *Am J Infect Control* 1984;12:247-248.
- ³²⁹ Doebbeling BN, Stanley GL, Sheetz CT, et al. Comparative efficacy of alternative handwashing agents in reducing nosocomial infections in 386 INFECTION CONTROL AND HOSPITAL EPIDEMIOLOGY May 2003 intensive care units. *N Engl J Med* 1992;327:88-93.
- ³³⁰ Ehrenkranz NJ, Alfonso BC. Failure of bland soap handwash to prevent hand transfer of patient bacteria to urethral catheters. *Infect Control Hosp Epidemiol* 1991;12:654-662.
- ³³¹ Larson E. A casual link between handwashing and risk of infection? Examination of the evidence. *Infect Control Hosp Epidemiol* 1988;9:28- 36.
- ³³² Larson EL, Eke PI, Laughon BE. Efficacy of alcohol-based hand rinses under frequent-use conditions. *Antimicrob Agents Chemother* 1986;30:542-544.
- ³³³ Larson EL, Aiello AE, Bastyr J, et al. Assessment of two hand hygiene regimens for intensive care unit personnel. *Crit Care Med* 2001;29:944- 951.
- ³³⁴ Boyce JM. Scientific basis for handwashing with alcohol and other waterless antiseptic agents. In: Rutala WA, ed. *Disinfection, Sterilization and Antisepsis: Principles and Practices in Healthcare Facilities*. Washington, DC: Association for Professionals in Infection Control and Epidemiology; 2001:140-151.
- ³³⁵ Widmer AF. Replace handwashing with use of a waterless alcohol rub? *Clin Infect Dis* 2000;31:136-143.
- ³³⁶ Maury E, Alzieu M, Baudel JL, Haram N. Availability of an alcohol solution can improve hand disinfection compliance in an intensive care unit. *Am J Respir Crit Care Med* 2000;162:324-327.

Appendix B

- ³⁵⁷ Boyce JM, Kelliher S, Vallande N. Skin irritation and dryness associated with two hand hygiene regimens: soap and water handwashing versus hand antiseptics with an alcoholic hand gel. *Infect Control Hosp Epidemiol* 2000;21:442-448.
- ³⁵⁸ Bischoff WE, Reynolds TM, Sessler CN, Edmond MD, Wenzel RP. Handwashing compliance by healthcare workers: the impact of introducing an accessible alcohol-based hand antiseptic. *Arch Intern Med* 2000;160:1017-1021.
- ³⁵⁹ Walsh B, Blakemore PH, Drubu YJ. The effect of handcream on the antibacterial activity of chlorhexidine gluconate. *J Hosp Infect* 1987;9:30-33.
- ³⁴⁰ Berndt U, Wigger-Alberti W, Gabard B, Elsner P. Efficacy of a barrier cream and its vehicle as protective measures against occupational irritant contact dermatitis. *Contact Dermatitis* 2000;42:77-80.
- ³⁴¹ McCormick RD, Buchman TL, Maki D. Double-blind, randomized trial of scheduled use of a novel barrier cream and an oil-containing lotion for protecting the hands of health care workers. *Am J Infect Control* 2000;28:302-310.
- ³⁴² Heeg P. Does hand care ruin hand disinfection? *J Hosp Infect* 2001;48(suppl A):S37-S39.
- ³⁴³ Dharan S, Hugonnet S, Pittet D. Evaluation of interference of a hand care cream with alcohol-based hand disinfection. *Occup Environ Med* 2001;49:81-84.
- ³⁴⁴ Mayer JA, Dubbert PM, Miller M, Burkett PA, Chapman S. Increasing handwashing in an intensive care unit. *Infect Control* 1986;7:259-262.
- ³⁴⁵ Quale J, Landman D, Atwood E, et al. Experience with a hospital-wide outbreak of vancomycin-resistant enterococci. *Am J Infect Control* 1996;24:372-379.
- ³⁴⁶ Muto CA, Durbin IJ, Alexander CH, Karchmer TB, Farr BM. Frequency of community acquired MRSA at a university hospital. Presented at the 35th Annual Meeting of the Infectious Diseases Society of America; September 13-16, 1997; San Francisco, CA. Abstract 347:135.
- ³⁴⁷ Harbarth S, Samore MH, Lichtenberg D, Carmeli Y. Prolonged antibiotic prophylaxis after cardiovascular surgery and its effect on surgical site infections and antimicrobial resistance. *Circulation* 2000;10:2916-2921.
- ³⁴⁸ Gross R, Kinky DE, Weiner M, Morgan AS, Gibson GA, Fishman NO. A randomized controlled trial of a comprehensive antimicrobial management program. *Infect Control Hosp Epidemiol* 2000;21:87-88.
- ³⁴⁹ Gross R, Morgan AS, Kinky DE, Weiner M, Gibson GA, Fishman NO. Impact of a hospital-based antimicrobial management program on clinical and economic outcomes. *Clin Infect Dis* 2001;33:289-295.
- ³⁵⁰ Evans RS, Pestonik SL, Classen DC, et al. A computer-assisted management program for antibiotics and other anti-infective agents. *N Engl J Med* 1998;338:232-238.
- ³⁵¹ May AK, Melton SM, McGwin G, Cross JM, Moser SA, Rue LW. Reduction of vancomycin-resistant enterococcal infections by limitation of broad-spectrum cephalosporin use in a trauma and burn intensive care unit. *Shock* 2000;14:259-264.
- ³⁵² Smith DW. Decreased antimicrobial resistance after changes in antibiotic use. *Pharmacotherapy* 1999;19:129S-132S.
- ³⁵³ Walker ES, Vasquez JE, Dula R, Bullock H, Sarubbi FA. Mupirocin-resistant, methicillin-resistant *Staphylococcus aureus*: does mupirocin remain effective? *Infect Control Hosp Epidemiol* 2003;24:342-346.

Appendix C

The Institute for Healthcare Improvement's 100,000 Lives Campaign



The Institute for Healthcare Improvement aims to enlist thousands of hospitals across the country in its 100,000 Lives Campaign. The goal is to save 100,000 lives a year by reducing nosocomial infections and medical errors. Three of the six elements of that campaign call on hospitals to implement procedures proven effective in reducing surgical site infections, central line infections, and ventilator-associated pneumonia. The three elements are summarized here. For the complete “How-To-Guide” to preventing these infections, consult the Institute’s Web site at: <http://www.ihl.org/IHI/Programs/Campaign>.

I. The Four Key Components of Preventing Ventilator-Associated Pneumonia:

1. Elevation of the head of the bed to between 30 and 45 degrees
2. Daily “sedation vacation” and daily assessment of the readiness to extubate
3. Peptic ulcer disease prophylaxis
4. Deep venous thrombosis prophylaxis (unless contraindicated)

II. The Four Key Components for Preventing Surgical Site Infections:

1. Appropriate use of antibiotics, including administering antibiotics within one hour before surgical incision, selecting an antibiotic consistent with national guidelines, and discontinuing prophylactic antibiotics within 24 hours after surgery
2. No shaving. Appropriate hair removal, if necessary, with clippers or a depilatory, but not with a razor
3. Monitor and maintain patient’s glucose levels after surgery, particularly for cardiovascular surgery patients.
4. Keep patients’ body temperatures at normal levels during and after surgery, especially colorectal surgery, with warmed IV fluids, warming blankets, hats and booties, and other means.

Appendix C

III. The Five Key Components of Preventing Catheter-Related Bloodstream Infections:

1. Appropriate hand hygiene, including cleaning hands before and after palpating catheter insertion sites, before and after inserting, replacing, accessing, repairing or dressing an intravascular catheter, whenever hands are soiled or contaminated, before and after removing gloves, etc.
2. Maximal barrier protection—meaning wearing a cap, mask, sterile gown, and gloves—when placing or assisting in the placement of a central line, and ensuring that the patient is covered head to toe in a sterile drape with one small opening for the site of insertion
3. Chlorhexidine skin antisepsis before insertion
4. Optimal catheter site selection, with the subclavian vein as the preferred site instead of the jugular or femoral sites for non-tunneled catheters in adult patients
5. Daily review of central line necessity to prevent unnecessary, prolonged use

About the Author

Dr. Betsy McCaughey is a health policy expert who has won many prizes for her writings, lectures widely, and appears frequently on television and radio. One year ago she launched a nationwide crusade to stop hospital infection deaths. She is founder and Chairman of the Committee to Reduce Infection Deaths (<http://www.hospitalinfection.org>).

Dr. McCaughey's research on how to prevent infection deaths has been featured on *Good Morning America*, the *CBS Morning Show*, ABC's *20/20*, and many other national programs. She has also appeared on Fox News Network's *Hannity & Colmes*, *The O'Reilly Factor*, CNN's *Talk Back Live*, and numerous radio programs.

Dr. McCaughey is the author of over one hundred scholarly or popular articles on health policy, which have appeared in *The New York Times*, *The Wall Street Journal*, *The New York Sun*, *New Republic*, *Policy Review*, *Forbes Magazine*, *New York Law Journal*, *Los Angeles Times*, *U.S. News & World Report*, and many other national publications.

She has produced prize-winning studies while at two think tanks, the Manhattan Institute (1993-94) and later the Hudson Institute (1999-2001). Dr. McCaughey's 1994 article on the dangers of the Clinton health plan received the National Magazine Award for the best article in the nation on public policy, the H. L. Mencken Award and other prizes. As Lt. Governor of New York State (1994-98), she proposed health legislation that became models for legislation in other states and in Congress. She has also been honored by the American Society of Anesthesiologists for her writings in that field.

Prior to entering the health policy field, Dr. McCaughey taught and wrote about U.S. constitutional history. She is the author of two books, *From Loyalist to Founding Father* (Columbia University Press), winner of the Bancroft Dissertation Award, and *Government by Choice* (Basic Books). She also chaired a national commission on reforming the electoral college in 1992, wrote its report, *Electing the President*, and testified before Congress on the subject. She has taught at Vassar College (1977-78) and Columbia University (1979-83), and in 1989, she served as Guest Curator for the Bicentennial Exhibit and related events at the New York Historical Society.

She is also the proud mother of three daughters: Amanda (Yale class of 2001), Caroline (Brown class of 2003) and Diana (Manhattanville class of 2006).

Endnotes

- ¹ Robert Weinstein, "Nosocomial Infection Update," *Journal of Emerging Infectious Diseases*, vol.4, no. 3. The infection rate, between 5 and 6 hospital-acquired infections per 100 admissions, has been steady for over the last 28 years, according to Weinstein's research.
 - ² The Centers for Disease Control and Prevention previously estimated that 90,000 people die annually from infections they contract in U.S. hospitals, but the CDC press confirms that it is "working on a new number." *The Chicago Tribune* puts the death rate even higher, at 103,000. *The Chicago Tribune* examined hospital records, court records, and federal and state agency data pertaining to 5810 hospitals to reach its estimate. The CDC based their extrapolations on data voluntarily submitted by 315 U.S. hospitals: "How the Chicago Tribune Analyzed Infection Cases," *Chicago Tribune* (July 21,2002).
 - ³ The Centers for Disease Control and Prevention previously estimated that 90,000 people die annually from infections they contract in U.S. hospitals, but the CDC press confirms that it is "working on a new number." Interview with Nicole Coffin, spokesperson for the CDC, Infectious Diseases Division, October 10, 2002 (404-639-2888). *The Chicago Tribune* puts the death rate even higher, at 103,000. *The Chicago Tribune* examined hospital records, court records, and federal and state agency data pertaining to 5,810 hospitals to reach its estimate. The CDC based their extrapolations on data voluntarily submitted by 315 U.S. hospitals: "How the Chicago Tribune Analyzed Infection Cases," *Chicago Tribune* (July 21,2002). See also Carloarena, M.D., et.al., "Assessment of Handwashing Practices with Chemical and Microbiological Agents," *American Journal of Infection Control*, (October, 2002) vol. 30., no. 6.; Didier M. Pillet, M.D., "Improving Adherence to Hand Hygiene," *Journal of Emerging Infectious Diseases* vol. 4, no. 3.; Leela C. Blamit, E. Louise Teare, William W. Williams, Jeremy D. Tuite, "Eradication of Methicillin Resistant Staphylococcus Aureus by 'ring fencing' of elective orthopedic beds," *British Medical Journal* vol. 329 (July 17, 2004).
 - ⁴ Institute of Medicine, *Insuring America's Health* (Washington, D.C., National Academies Press, 2004) 46.
 - ⁵ The Centers for Disease Control and Prevention previously estimated that 90,000 people die annually from infections they contract in U.S. hospitals, but the CDC press confirms that it is "working on a new number." Interview with Nicole Coffin, spokesperson for the CDC, Infectious Diseases Division, October 10, 2002 (404-639-2888). *The Chicago Tribune* puts the death rate even higher, at 103,000. *The Chicago Tribune* examined hospital records, court records, and federal and state agency data pertaining to 5,810 hospitals to reach its estimate. The CDC based their extrapolations on data voluntarily submitted by 315 U.S. hospitals: "How the Chicago Tribune Analyzed Infection Cases," *Chicago Tribune* (July 21,2002). See also Carloarena, M.D., et.al., "Assessment of Handwashing Practices with Chemical and Microbiological Agents," *American Journal of Infection Control*, (October, 2002) vol. 30., no. 6.
 - ⁶ See the "Cost of Hospital Infections" section of this booklet.
 - ⁷ Institute of Medicine, *Insuring America's Health* (Washington, D.C., National Academies Press, 2004) 46.
 - ⁸ Infectious Disease Society of America, *Bad Bugs, No Drugs*, (July, 2004) p. 3. Data are from the Centers for Disease Control and Prevention. National Nosocomial Infections Surveillance (NNIS) system report, data summary from January 1992 to June 2003, issued August 2003.
 - ⁹ "SHEA Guideline for Preventing Nosocomial Transmission of Multidrug-Resistant Strains of *Staphylococcus aureus* and *Enterococcus*," *Infection Control and Hospital Epidemiology*, 2003; vol. 24; pp. 362-386. See p.362.
 - ¹⁰ Infectious Disease Society of America, *Bad Bugs, No Drugs*, Table 3, p. 11. Data are drawn from the CDC National Nosocomial Infections Surveillance System.
 - ¹¹ "SHEA Guideline for Preventing Nosocomial Transmission of Multidrug-Resistant Strains of *Staphylococcus aureus* and *Enterococcus*," *Infection Control and Hospital Epidemiology*. See p. 364.
-
-

-
- ¹² Cassandra D. Salgado and Barry M. Farr, "MRSA and VRE: Preventing Patient-to-Patient Spread," *Infections in Medicine* (2003) vol: 20; pp. 192-200; "Epidemiology of nosocomial infections caused by *methicillin-resistant Staphylococcus aureus*," *Annals of Internal Medicine* (1982), vol. 97: pp. 309-317.
- ¹³ Pittsburgh Regional Healthcare Initiative, *PRHI in Progress: 2004 in Review* (January 2005) p. 5.
- ¹⁴ Telephone interview with Dr. Carlene Muto, University of Pittsburgh Medical Center-Presbyterian Hospital, July 15, 2005 (412-629-2566).
- ¹⁵ "Control of *vancomycin-resistant Enterococcus* in a region," *New England Journal of Medicine*, 2001:vol. 344: pp.1427-1433; also "MRSA and VRE: Preventing Patient-to-Patient Spread," *Infections in Medicine*, 2003: vol. 20; pp. 194-200.
- ¹⁶ Hand Hygiene in Healthcare Settings Core, slide 3 of a slide presentation dated June 6, 2003, and currently available on the CDC Web site at http://cdc.gov/handhygiene/download/hand_hygiene_core.ppt; See also: Didier M. Pillet, M.D., "Improving Adherence to Hand Hygiene," *Journal of Emerging Infectious Diseases* vol. 4, no. 3.; Leela C. Blant, E. Louise Teare, William W. Williams, Jeremy D. Tuite, "Eradication of *Methicillin Resistant Staphylococcus Aureus* by 'ring fencing' of elective orthopedic beds," *British Medical Journal* vol. 329 (July 17, 2004).
- ¹⁷ MRSA contaminates patient care rooms 69 percent to 73 percent of the time. "Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: possible infection control implications," *Infection Control and Hospital Epidemiology*, 1997: vol. 9, pp. 622-7.
- ¹⁸ "SHEA Guideline for Preventing Nosocomial Transmission of Multidrug-Resistant Strains of *Staphylococcus aureus* and *Enterococcus*," *Infection Control and Hospital Epidemiology*. See p.374.
- ¹⁹ Salgado and Farr, "MRSA and VRE," pp. 192-200
- ²⁰ *Ibid.*; See also: "Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: possible infection control implications," *Infection Control and Hospital Epidemiology*(1999) v01.9: 622-627; "Isolation gowns prevent health care workers (HCWs) from contaminating their clothing, and possibly their hands with methicillin-resistant *Staphylococcus aureus* (MRSA) and resistant enterococci, Presented to the Eighth Annual Meeting of the Society for Healthcare Epidemiology of America , April 5-7, 1998, Orlando, Florida, Abstract No. S74:52, cited in "SHEA Guideline for Preventing Nosocomial Transmission of Multidrug-Resistant Strains of *Staphylococcus aureus* and *Enterococcus*," *Infection Control and Hospital Epidemiology* (May, 2003) vol. 24: JEAS.
- ²¹ Separate discussions with two Mt. Sinai physicians: Dr. Jennifer Glueck, 212-410-4068, resident in OB/GYN, and Dr. Douglas Dietrich, Vice Chairman of the Department of Medicine at 212-241-7270.
- ²² The American Medical Association recommends it. "MRSA and VRE: Preventing Patient to Patient Spread."
- ²³ Telephone interview with Dr. Carlene Muto, May 19, 2005 at 412-629-2566.
- ²⁴ Jernigan, J.A., Titus, M.G., Farr B.M., Groschel, D.H.M., Getchell-White, S.I., "Effectiveness of Contact Isolation during a Hospital Outbreak of Methicillin-Resistant *Staphylococcus aureus*," *American Journal of Epidemiology* (1996) vol. 146, p. 496-504. Cited in "SHEA Guideline for Preventing Nosocomial Transmission of Multidrug-Resistant Strains of *Staphylococcus aureus* and *Enterococcus*," footnote 181.
- ²⁵ Presentation by Dr. Richard Shannon, Chairman of the Department of Medicine, Allegheny Hospital at the March 11, 2005 Forum, slide no. 13, "The Compelling Case for Reducing Hospital Infection," Harvard Club of New York City.
- ²⁶ "Cost-Effective Eradication of an Outbreak of Methicillin-Resistant *Staphylococcus aureus* in a Community Teaching Hospital," *Infection Control and Hospital Epidemiology* (1988) vol: 9: pp. 255-260.
- ²⁷ "The Use of Economic Modeling to Determine the Hospital Costs Associated with Nosocomial Infections," *Clinical Infectious Diseases* (2003) vol. 36: pp. 1424-1432. This study puts the average cost of hospital infection at \$15,275. "A Systematic Audit of Economic Evidence Linking Nosocomial Infections and Infection Control Interventions, 1990-2000," *American Journal of Infection Control*,(2002), vol 30: pp. 145-52. This study
-

estimates the average cost of infection to be \$13,973. Both studies look at incremental hospital costs only. The national cost was reached by multiplying 2 million infections by \$15,000 per infection.

- ²⁸ "The Impact of Antimicrobial Resistance on Health and Economic Outcomes," *Clinical Infectious Diseases*, 2003; vol. 36: pp. 1433-1437. The investigators found that antibiotic resistance increases length of stay, mortality rates, and costs.
- ²⁹ Rebecca R. Roberts, R. Douglas Scott II, Ralph Cordell, Steven L. Solomon, Lynn Steele, Linda M. Kampe, William E. Trick, and Robert A. Weinstein, "The Use of Economic Modeling to Determine the Hospital Costs Associated with Nosocomial Infections," *Clinical Infectious Diseases* (2003) vol. 36: 1424-1432. The figure \$15,275 refers to excess hospital costs, not other patient care costs or economic losses incurred by the patient as a result of prolonged illness. The figure \$15,275 is similar to findings from other analyses of average hospital care costs due to infection. MedMined, Inc. analyzed the costs of infection using data from 50 hospital clients. It found that patients contracting infections in the hospital stayed an average of eight days extra and incurred \$14,000 in additional hospital costs. See: John Morrissey, "Debugging Hospitals," *Modern Healthcare* (April 26, 2004): 30.
- ³⁰ Nancy M. Kane, Richard B. Siegrist, Jr., "Understanding Rising Hospital Inpatient Costs: Key Components of Cost and the Impact of Poor Quality," (August 12, 2002), unpublished manuscript. The findings are based on Medicare Cost Reports (1999-2000) and Uniform Hospital Discharge Data Set (2000) for all acute general hospitals in ten states: Massachusetts, New York, Virginia, Florida, Texas, Illinois, Iowa, California, Washington, and Colorado. Kane and Siegrist are adjunct faculty at the Harvard School of Public Health.
- ³¹ Ibid.
- ³² Institute for Healthcare Improvement, 100,000 Lives Campaign, How-to-Guide: Prevent Ventilator-Associated Pneumonia at www.ihp.org/IHI/Programs/Campaign.
- ³³ Pittsburgh Regional Healthcare Initiative, "PRHI Executive Summary," (January, 2005). Dr. Richard Shannon found that each central line-associated bloodstream infection added \$30,277 in variable costs during the hospital stay, for patients in the medical intensive care unit and coronary care unit at Allegheny Hospital, Pittsburgh.
- ³⁴ Gary A. Noskin; Robert J. Rubin; Jerome J. Schentag; Jan Kluytmans; Edwin C. Hedblom; Maartje Smulders; Elizabeth Lapetina; Eric Gemmen, "The Burden of *Staphylococcus aureus* Infections on Hospitals in the United States," *Archives of Internal Medicine* vol. 165 (2005): 1756-1761.
- ³⁵ Twenty-one states require hospitals to report "adverse events" including these serious infections. The best source for these state requirements is the National Academy for State Health Policy, Portland, Maine. At www.nashp.org
- ³⁶ The CDC's National Nosocomial Infections Surveillance System, including the guarantees of confidentiality to hospitals, is described at www.cdc.gov/ncidod/hip/SURVEILL/NNIS.HTM
- ³⁷ Mark Chassin, "Achieving and Sustaining Improved Quality: Lessons from New York State and Cardiac Surgery," *Health Affairs*, (July/August 2002) vol. 21, no. 4, 40-51.
- ³⁸ Ibid.
- ³⁹ Judith H. Hibbard, Jean Stockard, Martin Tusler, "Does Publicizing Hospital Performance Stimulate Quality Improvement Efforts," *Health Affairs* (March/ April 2003), vol. 22, no. 2, 84-94.
- ⁴⁰ For more information on the NNIS's methodology, consult: TC Horan & R. Gaynes, "Surveillance of nosocomial infections," *Hospital Epidemiology and Infection Control*, ed. CG. Mayhall (Philadelphia: Lippincott, Williams & Wilkins, 2004) 1659-1702.
- ⁴¹ J. Steinhauer, "Hospitals in City Faulted for Failing to Report Many Errors," *New York Times* (February 13, 2001), B1; see also Mark Chassin, "The Wrong Patient," *Annals of Internal Medicine* (June 4, 2002) vol. 136, no. 11 836-831.
- ⁴² "Data Show Scourge of Hospital Infections," *Washington Post*, (July 13, 2005); "Hospitals Battling Infections," *Washington Post*, (July 14, 2005).
- ⁴³ Telephone interview with Dr. Frank Lowey, Professor of Microbiology, Columbia College of Physicians and Surgeons, July 15, 2005.
- ⁴⁴ The CDC and the American Association of Medical Colleges are working on a Web-based curriculum that focuses largely on this issue.
- ⁴⁵ Based on an interview with Dr. Carlene Muto by telephone, July 15, 2005.
-
-

-
- ⁴⁶ Dr. Richard P. Shannon, PowerPoint presentations, December 7, 2004 and March 11, 2005, Harvard Club of New York City; March 11, 2005, Harvard Club of New York City; "PRHI in Progress: 2004 in Review," published by the Pittsburgh Regional Healthcare Initiative.
- ⁴⁷ Dr. Richard Shannon, PowerPoint presentations, December 7, 2004 and March 11, 2005.
- ⁴⁸ *Ibid.*
- ⁴⁹ "Reducing hospital-acquired infections," *Ivanhoe Broadcast News*, (9-25-2004) at www.news8austin.com; "Simple Intervention Nearly Eliminates Catheter-related Bloodstream Infections," (12-09-2004) at www.sciencedaily.com
- ⁵⁰ Joe Fahy, "Hospitals Able to Cut Some Kinds of Infections," *Pittsburgh Post-Gazette*, July 15, 2005, A1.
- ⁵¹ Based on an interview with Dr. Barry Farr by telephone, July 21, 2005.
- ⁵² "Impact of Ring Wearing on Hand Contamination and Comparison of Hand Hygiene Agents in a Hospital," *Clinical Infectious Diseases* 36 (2003): 1383-1390.
- ⁵³ Nearly three quarters of patients' rooms are contaminated with MRSA, and 69 percent with VRE, studies show. In one study, 42 percent of gloves worn by hospital personnel who had no direct patient contact but who touched contaminated surfaces became contaminated. "Environmental contamination due to *methicillin-resistant Staphylococcus aureus*: possible infection control implications," *Infection Control and Hospital Epidemiology* 9 (1997): 622-627. A Consensus Statement by a multidisciplinary group of experts asked by the American Medical Association to provide guidelines for infection control cautions that "In some cases caregivers actually go from patient to patient without changing their gloves, apparently confusing self-protection" with patient protection. "Strategies to Prevent and Control the Emergence and Spread of Antimicrobial-Resistant Microorganism in Hospitals," *Journal of the American Medical Association* 275 (1996): 234-240.
- ⁵⁴ Routine disinfection of stethoscopes between patients is recommended by the American Medical Association. "MRSA and VRE: Preventing Patient-to-Patient Spread," *Infections in Medicine* 20 (2003):194-200; "The Stethoscope: a powerful source of nosocomial infection?" *Archives of Internal Medicine*,157 (1997): 786-90; "Contamination of gowns, gloves, and stethoscopes with vancomycin-resistant Enterococci," *Infection Control and Hospital Epidemiology* 9 (2001): 560-564; "Recovery of vancomycin-resistant Enterococci on fingertips and environmental surfaces," *Infection Control and Hospital Epidemiology* 12 (1996): 770-772.
- ⁵⁵ "Eradication of *methicillin-resistant Staphylococcus aureus* by 'ring fencing' of elective orthopaedic beds," *British Medical Journal*,p. 149-51. Visitors who sit on a chair or lean on a cabinet, then sit on the bed are transferring bacteria to the patient's bedclothes. "MRSA and VRE," *Infections in Medicine*, p. 194-200.
- ⁵⁶ The Institute for Healthcare Improvement guidelines for improving infection prevention state that: "Administration of prophylactic antibiotics beginning 0 to 1 hour prior to surgical incision decreases the risk of surgical infection. www.imi.org/IHI/Topics/PatientSafety/SurgicalSiteInfections/ImprovementStories accessed 10-14-02.
- ⁵⁷ *Ibid.*, the Institute for Healthcare Improvement Guidelines for improving infection state that "surgical patients with core temperatures greater than 36 degrees C./ 98.6 degrees F are less likely to get an infection."
- ⁵⁸ *Ibid.*, the Institute for Healthcare Improvement states that "clipping instead of shaving results in decreased infection rates," and recommends that patients be told "not to shave the surgical site for 72 hours prior to surgery."
- ⁵⁹ "Operating Room Environment" *Clinical Orthopaedics and Related Research* 369 (1999): 103-109.
- ⁶⁰ Pittsburgh Regional Healthcare Initiative, "PRHI Executive Summary," (June, 2005).
- ⁶¹ Urinary tract infections are the most common hospital-acquired infections. Limiting their use and duration reduces risk of infection. "Catheter Associated Urinary Tract Infections in Neurology and Neurosurgical Units," *Journal of Infection* 44 (2002): 171-175.
- ⁶² These are the types of infections that the Centers for Disease Control and Prevention recommend in a public reporting system. However, lawmakers could consider adding ventilator-related pneumonia, but these are more difficult to detect accurately, or urinary tract infections, which are common but less costly and deadly than other infections.
-